

NEW EXAMPLES OF OBSTRUCTIONS TO NON-NEGATIVE SECTIONAL CURVATURES IN COHOMOGENEITY ONE MANIFOLDS

CHENXU HE

ABSTRACT. K. Grove, L. Verdiani, B. Wilking and W. Ziller gave the first examples of cohomogeneity one manifolds which do not carry invariant metrics with non-negative sectional curvatures. In this paper we generalize their results to a larger family. We also classified all class one representations for a pair (G, H) with G/H some sphere, which are used to construct the examples.

1. INTRODUCTION

Non-negatively curved Riemannian manifolds have been of interest from the beginning of global Riemannian geometry. Most examples are obtained via product and quotient constructions, starting from compact Lie groups. These include all homogeneous spaces and biquotients. Using a gluing method, J. Cheeger constructed non-negatively curved metrics on the connect sum of any two rank one symmetric spaces, see [Ch]. A breakthrough came with K. Grove and W. Ziller's generalization of this gluing method to the class of cohomogeneity one manifolds.

A manifold M is called a *cohomogeneity one manifold* if there exists a compact Lie group G acting on M by isometries and the cohomogeneity of the action, defined as $\text{cohom}(M, G) = \dim(M/G)$, is equal to 1. Since the orbit space is one dimensional, it is either a circle or a closed interval I . In the former case, M always carries a G invariant metric with non-negative sectional curvature. In the latter one, there are precisely two singular orbits $B_{\pm}$ with isotropy subgroups $K^{\pm}$ corresponding to the endpoints of I , a minimal geodesic between the singular orbits, and principal orbits corresponding to the interior points with isotropy subgroup H . By the slice theorem, $K^{\pm}/H$ are spheres $\mathbb{S}^{l_{\pm}-1}$ ($l_{\pm} \geq 1$), and M can be reconstructed by gluing two disk bundles along a principal orbit as follows

$$(1.1) \quad M = G \times_{K^-} \mathbb{D}^{l_-} \cup_{G/H} G \times_{K^+} \mathbb{D}^{l_+},$$

where $\mathbb{D}^{l_{\pm}}$ is the normal disk to $B_{\pm}$. Therefore we can identify M with the groups $H \subset \{K^-, K^+\} \subset G$ by the gluing construction (1.1).

K. Grove and W. Ziller showed that any cohomogeneity one manifold with codimension two singular orbits admits a non-negatively curved metric [GZ1]. In the same paper, it was

conjectured that any cohomogeneity one manifold admits a non-negatively curved metric. This turns out to be false. The first examples of an obstruction were discovered by K.Grove, L.Verdiani, B.Wilking and W.Ziller in [GVWZ]. The most interesting examples are the higher dimensional Kervaire spheres (of dimension 9 and up) which are the only exotic spheres that can carry a cohomogeneity one action, see [St].

In light of the construction of examples with non-negative sectional curvature and the examples of obstructions to non-negatively curved metrics, it is important to answer the question "How large is the class of cohomogeneity one manifolds that admit a non-negatively curved metric?" which was raised by W.Ziller in [Zi].

In this paper, we generalize the examples in [GVWZ] to a larger family:

Theorem. *Let $K'/H' = \mathbb{S}^k$ with $k \geq 2$ and $\rho : K' \longrightarrow SO(m)$ be a faithful irreducible representation, which is not the one of K' on $\mathbb{R}^{k+1}$, and such that $\rho(H') \subset SO(m-1)$. For any integer $n \geq m+2$, set $G = SO(n)$ and*

$$\begin{aligned} K^- &= \rho(K') \times SO(n-m) \subset SO(m) \times SO(n-m) \subset SO(n) \\ K^+ &= \rho(H') \times SO(n-m+1) \subset SO(m-1) \times SO(n-m+1) \subset SO(n) \\ H &= \rho(H') \times SO(n-m) \subset SO(n), \end{aligned}$$

then the cohomogeneity one manifold M defined by the groups $H \subset \{K^-, K^+\} \subset G$ does not admit a G invariant metric with non-negative sectional curvature.

In [GVWZ], the theorem was proved under the additional assumptions that the slice representation of K' is not contained in the symmetric square $\text{Sym}^2 \rho$ and $\rho(K')$ does not act transitively on the sphere $\mathbb{S}^{m-1} = SO(m)/SO(m-1)$.

This theorem is optimal in the sense that if ρ is the representation of K' on $\mathbb{R}^{k+1}$, then the manifold M does admit an invariant non-negatively curved metric, since it is diffeomorphic to the homogeneous space $SO(n+1)/(\rho(K') \times SO(n-m+1))$ endowed with the cohomogeneity one action of $SO(n) \subset SO(n+1)$. On the other hand, if $n = m+1$ one obtains an interesting cohomogeneity one manifold where it is not yet known whether that manifold carries a non-negatively curved invariant metric or not.

The theorem can be extended to the case where ρ is not necessarily irreducible, see Theorem 3.1. A representation $\rho : K' \longrightarrow SO(m)$ with $\rho(H') \subset SO(m-1)$ is called a *class one* representation, see Definition 2.7. Many class one representations are not faithful, see Proposition A.19 and Table 2, so they are excluded by the faithfulness requirement in [GVWZ]. However Theorem 3.1 allows non-faithful class one representation as a subrepresentation of ρ and it gives us many more examples, see Section 3.

Similar results also hold if ρ is a complex or quaternionic representation, i.e., $G = U(n)$ or $Sp(n)$ which are stated in the Theorem 6.1 and Theorem 6.6.

In the above theorem, Theorem 3.1, Theorem 6.1 and Theorem 6.6, the group K' does not need to be connected. But the corresponding groups $K^\pm$ on the universal cover $\widetilde{M}$ will be connected. Therefore in the rest of the paper, we can assume that all groups are connected.

The paper is organized as follows. In Section 2, we briefly recall basic properties of cohomogeneity one manifolds, the generalized Weyl group and metric properties. Also this section includes an introduction to class one representations. A more detailed discussion of such representations is given in Appendix A. In Section 3 we prove some basic properties of the new examples which will be used in the proof of the orthogonal case. In Section 4, we study the consequences of the non-negativity assumption on the metrics. Wilking's rigidity theorem for non-negatively curved manifolds[Wi1] plays an important role. Section 5 is devoted to the proof in the orthogonal case. Using properties of the metrics developed in the previous sections, we draw a contradiction by looking at sectional curvatures of 2 distinct classes of 2-planes. In the last section, we sketch an outline of the proofs in the case where $G = U(n)$ or $Sp(n)$.


Acknowledgements: The paper is part of the author's Ph.D. thesis at University of Pennsylvania. The author wants to thank his advisor, Prof. Wolfgang Ziller, for his generous supports and great patience, and to Prof. Kristopher Tapp for valuable discussions on Wilking's rigidity results.

2. PRELIMINARIES

In this section, we recall some basic and well-known facts about cohomogeneity one manifolds. For more detail, we refer to, for example, [AA] and [GWZ].

As mentioned already, there are precisely two non-principal orbits $B_\pm$ in a simply connected cohomogeneity one manifold. Suppose M is endowed with an invariant metric g and the distance between the two non-principal orbits is L . Let $c(t)$, $t \in \mathbb{R}$ be a geodesic minimizing the distance with $c(0) = p_- \in B_-$ and $c(L) = p_+ \in B_+$. The isotropy subgroups at $p_\pm$ are denoted by $K^\pm$ and the principal isotropy subgroup at any point $c(t)$, $t \in (0, L)$, is

denoted by H . We can draw the following group diagram for the manifold M :



The group diagram $H \subset \{K^-, K^+\} \subset G$ is not uniquely determined by the manifold since one can switch K^- with K^+ , change g to another invariant metric and choose another minimal geodesic $c(t)$.

Definition 2.1. Two group diagrams are called *equivalent* if they determine the same cohomogeneity one manifold up to equivariant diffeomorphism.

The following lemma characterizes which two group diagrams are equivalent, see in [GWZ].

Lemma 2.2. *Two group diagrams $H \subset \{K^-, K^+\} \subset G$ and $\tilde{H} \subset \{\tilde{K}^-, \tilde{K}^+\} \subset G$ are equivalent if and only if after possibly switching the roles of K^- and K^+ , the following holds: There exist elements $b \in G$ and $a \in N(H)_0$, where $N(H)_0$ is the identity component of the normalizer of H , with $\tilde{K}^- = bK^-b^{-1}$, $\tilde{H} = bHb^{-1}$, and $\tilde{K}^+ = abK^+b^{-1}a^{-1}$.*

Remark 2.3. If $c(t)$ is the minimal normal geodesic between the two singular orbits, then $b \star c(t)$ is another minimal geodesic between $B_\pm$ and the associated group diagram is obtained by conjugating all isotropy groups by the element b . We can assume that $b \in N(H) \cap N(K^-)$ in order to fix H and K^- . Conjugation by an element a as in the above lemma usually corresponds to changing the invariant metric on the manifold.

Let $C \subset M$ be the image of the minimal geodesic $c(t)$. Then the Weyl group $\mathcal{W}$ of the G -action on M is by definition the stablizer of C modulo its kernel H . $\mathcal{W}$ is characterized in the following proposition.

Proposition 2.4. *The Weyl group $\mathcal{W}$ of a cohomogeneity one manifold is a dihedral subgroup of $N(H)/H$. It is generated by involutions $w_\pm \in (N(H) \cap K^\pm)/H$ and $C/\mathcal{W} = M/G = [0, L]$. Each of these involutions can be represented as an element $a \in K^\pm \cap H$ such that a^2 but not a lies in H .*

Using the group action, the invariant metric is determined by its restriction to the minimal geodesic $c(t)$. Suppose $c(t)$ is parameterized by arc length, i.e., $T = \frac{d}{dt}$ has length 1, then

we can write g as

$$g = dt^2 + g_t,$$

and $\{g_t\}_{t \in [0, L]}$ is a one-parameter family of homogeneous metrics on the orbits $M_t = G.c(t)$.

Fix a bi-invariant inner product Q on the Lie algebra $\mathfrak{g}$ of G and let $\mathfrak{p} = \mathfrak{h}^\perp$ be the orthogonal complement of the Lie algebra $\mathfrak{h}$ of H . For each $X \in \mathfrak{p}$, let X^* be the Killing vector field generated by X along $c(t)$, i.e., $X^*(t) = \frac{d}{ds}|_{s=0} \exp(sX).c(t)$. For each $t \in (0, L)$, M_t is diffeomorphic to the homogeneous space G/H , and hence $T_{c(t)}M_t$ can be identified with $\mathfrak{p}$ by means of Killing vector fields as $X \mapsto X^*(t)$. Then g_t defines an inner product on $\mathfrak{p}$ which is invariant under the isotropy action of Ad_H . We set

$$(2.5) \quad g_t(X, Y) = g_t(X^*(t), Y^*(t)) = Q(P_t X, Y) \text{ for } X, Y \in \mathfrak{p},$$

where $P_t : \mathfrak{p} \longrightarrow \mathfrak{p}$ is a Q -symmetric Ad_H -equivariant endomorphism. The metric g is completely determined by the one parameter family $\{P_t\}$, $t \in [0, L]$, and at $t = 0$ and L , P_t should satisfy further conditions to guarantee smoothness of g .

On the other hand, each principal orbit M_t is a hypersurface in M with normal vector T . If $S_t X = S_t X^*(t) = -\nabla_{X^*} T$ is the shape operator of M_t at $c(t)$, we have

$$(2.6) \quad S_t = -\frac{1}{2} P_t^{-1} P'_t$$

in terms of P .

In the rest of this section, we give a short introduction to class one representations with more details in Appendix A. This particular class of representations is well studied, see, for examples, [VK] and [Wa].

First we recall:

Definition 2.7. A representation (μ, W) of a compact Lie group K is called a *real (complex or quaternionic) representation* if W is a vector space over $\mathbb{R}$ ($\mathbb{C}$ or $\mathbb{H}$).

Definition 2.8. Suppose W is an irreducible representation over the complex numbers, then W is called a *real representation* or of *real type* if it comes from a representation over reals by extension of scalars. It is of *quaternionic type* if it comes from a representation over quaternions by restriction of scalars. It is of *complex type* if it is neither real or quaternionic.

For any complex representation μ , let μ^* denote its complex conjugate. μ^* is equivalent to μ if μ is of type real or quaternionic and they are non-equivalent if μ is of complex type.

Suppose K is a compact connected Lie group, then the complexification of a real irreducible representation σ is one of the following

- (1) $\sigma_{\mathbb{C}} = \mu$, where μ is an irreducible representation of real type,

- (2) $\sigma_{\mathbb{C}} = \mu \oplus \mu^*$, where μ is irreducible and of complex type,
- (3) $\sigma_{\mathbb{C}} = \mu \oplus \mu$, where μ is irreducible and of quaternionic type.

In class (2) and (3), we often write $\sigma = [\mu]_{\mathbb{R}}$.

Conversely, suppose μ is a complex irreducible representation of K with degree n . If μ is of real type, then there exists a real vector space $\mathbb{R}^n \subset \mathbb{C}^n$ which is $\mu(K)$ invariant. Let σ be the restriction of μ on $\mathbb{R}^n$, then σ is a real irreducible representation with $\sigma_{\mathbb{C}} = \mu$. If μ is not of real type, then we identify $\mathbb{C}^n$ with $\mathbb{R}^{2n}$ and forget the complex structure on it. Thus $\mathbb{R}^{2n}$ is invariant under the $\mu(K)$ action and it gives us a real irreducible representation. We denote this representation by σ and then $\sigma_{\mathbb{C}}$ is either $\mu \oplus \mu^*$ or $\mu \oplus \mu$ depending on the type of μ .

A quaternionic irreducible representation is obtained by extending the scalar field to the quaternions for a complex irreducible representation and the converse also holds.

Given an n dimensional real irreducible representation (σ, W) of a compact Lie group K , we briefly discuss the classification of the equivariant endomorphisms, i.e., the endomorphism $f : W \longrightarrow W$ with $f(\sigma(g).v) = \sigma(g).f(v)$ for any $g \in K$ and $v \in W$.

If σ is in class (1), then from Schur's lemma, $f = a\text{Id}$ for some constant $a \in \mathbb{R}$ and Id is the identity map of W .

If σ is in class (2), then W has an orthonormal basis such that f is in the following form under that basis

$$(2.9) \quad f = \begin{pmatrix} a_0 I_m & -a_1 I_m \\ a_1 I_m & a_0 I_m \end{pmatrix},$$

where a_0, a_1 are constants in $\mathbb{R}$, $n = 2m$ and I_m is the $m \times m$ identity matrix.

If σ is in class (3), then there exists an orthonormal basis of W such that f has the following form under that basis

$$(2.10) \quad f = \begin{pmatrix} a_0 I_m & -a_1 I_m & -a_2 I_m & -a_3 I_m \\ a_1 I_m & a_0 I_m & a_3 I_m & -a_2 I_m \\ a_2 I_m & -a_3 I_m & a_0 I_m & a_1 I_m \\ a_3 I_m & a_2 I_m & -a_1 I_m & a_0 I_m \end{pmatrix},$$

where $n = 4m$, a_0, a_1, a_2 and a_3 are constants in $\mathbb{R}$.

Definition 2.11. A pair (K, H) of compact Lie groups with $H \subset K$ and $K/H = \mathbb{S}^k$ ($k \geq 2$) is called a *spherical pair*.

If we assume that K is connected, the image of μ will be a closed subgroup of $SO(l)$, $U(l)$ or $Sp(l)$ if μ is over $\mathbb{R}$, $\mathbb{C}$ or $\mathbb{H}$.

For each group pair (K, H) with $H \subset K$ a closed subgroup, we have

Definition 2.12. A non-trivial irreducible representation (μ, W) of K is called a *class one representation of the pair (K, H)* if $\mu(H)$ fixes a nonzero vector $w_0 \in W$.

Remark 2.13. From Proposition A.19, the class one representations of spherical pairs are almost effective, i.e., the kernel of μ is discrete, except for the following cases:

- (1) $(K, H) = (U(n), U(n-1)_m)$ and $\mu = ae_1 - ae_n$ where a is a positive integer. The kernel is the diagonal embedded $U(1) \subset U(n)$.
- (2) $(Sp(n) \times Sp(1), Sp(n-1) \times Sp(1))$ and $\mu = a\varpi_2$ where a is a positive integer. The kernel contains the $Sp(1)$ factor.
- (3) $(Sp(n) \times U(1), Sp(n-1) \times U(1)_m)$ and $\mu = (a\varpi_1 + b\varpi_2) \otimes \text{id}$ where a, b are non-negative integers with $a + b \geq 1$ and id is the trivial representation of $U(1)$. The kernel contains the $U(1)$ factor.

More information on the non-trivial kernels for class one representations is given in Proposition A.19 and listed in Table 2.

In the case where $(K, H) = (SO(k+1), SO(k))$, let ϖ_1 be the highest weight of the standard representation ϱ_{k+1} on $\mathbb{R}^{k+1}$, then the class one representations over $\mathbb{R}$ are precisely those with the highest weights as $m\varpi_1$, $m = 1, 2, \dots$. These representation spaces can be realized as the space of homogeneous harmonic polynomials. Let $\{x_1, \dots, x_{k+1}\}$ be the basis of $\mathbb{R}^{k+1}$ and $SO(k+1)$ act by the matrix multiplication. Then an element $A \in SO(k+1)$ acts on a polynomial $f(x_1, \dots, x_{k+1})$ through the action on the variables, i.e., $(A.f)(x_1, \dots, x_{k+1}) = f(A^{-1}.x_1, \dots, A^{-1}.x_{k+1})$.

A polynomial f is called harmonic if $\Delta f = 0$ where $\Delta = \sum_{i=1}^{k+1} \frac{d^2}{dx_i^2}$. Let H_m be the space of homogeneous harmonic polynomials in $x_1, \dots, x_{k+1}$ of degree m , then the representation of $SO(k+1)$ on H_m has the highest weight $m\varpi_1$. All of them are of real type. If $k+1$ is odd, then the representation for any positive m is faithful. If $k+1$ is even, then the class one representation is faithful if and only if m is odd. As we will see that the class one representations of other spherical pairs (K, H) are the irreducible components when H_m is restricted to the subgroup K of $SO(k+1)$.

The real, complex and quaternionic class one representations of the spherical pairs are classified in Appendix A. Further properties are discussed there as well.

3. WEYL GROUP AND SMOOTHNESS

First let us state the most general result in the orthogonal case as a generalization of the theorem in the introduction using the concept of class one representation:

Theorem 3.1. *Let $K'/H' = \mathbb{S}^k$ with $k \geq 2$ and $\rho : K' \longrightarrow SO(m)$ be a faithful representation that contains a (not necessarily faithful) irreducible class one representation μ of the pair (K', H') such that one of the followings holds:*

- (1) $\deg \mu \geq k + 2$ if μ is of real type or the multiplicity of μ in ρ , denoted by $\text{mul}(\mu, \rho)$, is equal to 1;
- (2) $\deg \mu \geq 2(k + 2)$ if μ is of complex type and $\text{mul}(\mu, \rho) \geq 2$;
- (3) $\deg \mu \geq 4(k + 2)$ if μ is of quaternionic type and $\text{mul}(\mu, \rho) \geq 2$.

We assume that $n \geq m + 2$ if $\text{mul}(\mu, \rho) = 1$, and that $n \geq m + 3$ if $\text{mul}(\mu, \rho) \geq 2$. If we set $G = SO(n)$ and

$$\begin{aligned}
 K^- &= \rho(K') \times SO(n - m) \subset SO(m) \times SO(n - m) \subset SO(n) \\
 (3.2) \quad K^+ &= \rho(H') \times SO(n - m + 1) \subset SO(m - 1) \times SO(n - m + 1) \subset SO(n) \\
 H &= \rho(H') \times SO(n - m) \subset SO(n),
 \end{aligned}$$

then the cohomogeneity one manifold M defined by the groups $H \subset \{K^-, K^+\} \subset G$ does not admit a G invariant metric with non-negative sectional curvature.

Remark 3.3. Since we do not assume that μ is faithful, a non-faithful class one representation is allowed in this construction. In other words, if μ is a class one representation with $\deg \mu \geq k + 2$, we choose a representation τ with $\ker \tau \cap \ker \mu = \{1\}$. Then $\rho = \tau \oplus \mu$ satisfies the conditions in the theorem. For example, take $(K', H') = (SO(6), SO(5))$ with $k = 5$ and let μ be the 20 dimensional representation of $SO(6)$ with the highest weight $2\varpi_1$. To construct a cohomogeneity one manifold, we can choose $\tau = \varrho_6$ as the standard representation of $SO(6)$ which is faithful and let $\rho = \tau \oplus \mu$.

Remark 3.4. If ρ contains only one copy of μ or μ is of real type, then Table 3 in Proposition A.21 lists all real class one representations which have dimensions smaller than $k + 2$. We see that only the following representations are excluded by the assumption, $\deg \mu \geq k + 2$: the defining representation of K' on $\mathbb{R}^{k+1}$, the 9 dimensional representation ϱ_9 of the pair $(Spin(9), Spin(7))$, the 5 dimensional representation of $(Sp(2), Sp(1))$ and the 3 dimensional representations of $(SU(2), \{\text{Id}\})$ and $(U(2), U(1))$. All of these representations are not faithful, so one needs to add another representation τ with $\ker \tau \cap \ker \mu = \{\text{Id}\}$ to define a cohomogeneity one manifold. For such manifolds, we do not know if they admit an invariant metric with non-negative curvature.

Remark 3.5. If ρ contains more than one copy of μ and μ is not of real type, then the further restriction $\deg \mu \geq 2(k + 2)$ or $4(k + 2)$ excludes 8 more representations as listed in the last part of Table 3. Among them, the following representations are faithful: $\mu = [2\varpi_1]_{\mathbb{R}}$

for the pair $(SU(3), SU(2))$, $\mu = [\varpi_1 + \varpi_2]_{\mathbb{R}}$ for the pair $(Sp(2), Sp(1))$ and $\mu = [3\varpi_1]_{\mathbb{R}}$, $[5\varpi_1]_{\mathbb{R}}$ and $[7\varpi_1]_{\mathbb{R}}$ for the pair $(Sp(1), \{1\})$. The first one is of complex type and the other four are of quaternionic type. They can be used to construct cohomogeneity one manifolds without adding other representations, for example, take $(K', H') = (Sp(1), \{1\})$ and $\rho = [3\varpi_1]_{\mathbb{R}} \oplus [3\varpi_1]_{\mathbb{R}}$. Thus Theorem 3.1 does not give obstruction for such manifolds.

Remark 3.6. The lowest dimensional example of Theorem 3.1 is obtained as follows. Take $(K', H') = (SO(3), SO(2))$ with $K'/H' = \mathbb{S}^2$, then the lowest dimensional class one representation μ with $\deg > 3$ is the unique 5-dimensional representation of $SO(3)$ which is also faithful. If $m = 5$ and $n = 7$, the manifold M has dimension 20 and isotropy groups

$$\mu(SO(2)) \times SO(2) \subset \{\mu(SO(3)) \times SO(2), \mu(SO(2)) \times SO(3)\} \subset SO(7).$$

Notice that this example is already covered by Theorem 3.2 in [GVWZ].

We now describe the explicit embedding of the groups. $SO(m) \times SO(n-m)$ is embedded in $G = SO(n)$ block-wise, i.e., $SO(m)$ sits in the upper-left $m \times m$ -block and $SO(n-m)$ is in the lower-right block. By assumption, ρ is a faithful orthogonal representation of K' with representation space $V = \mathbb{R}^m$. Let $W_1, \dots, W_\alpha$ be invariant subspaces of V such that they are pairwise orthogonal, and the restrictions of $\rho(K')|_{W_i} = \mu$ are equivalent and irreducible. Let U be the orthogonal complement of $W = W_1 \oplus \dots \oplus W_\alpha$. According to the decomposition of V into invariant spaces, we can write ρ as

$$(3.7) \quad \rho = \tau \oplus \mu \oplus \dots \oplus \mu,$$

where τ is restriction of ρ to U and μ is the class one representation of (K', H') . Furthermore, μ is not a subrepresentation of τ . Suppose $r = \dim U$, $l = \dim W_i = \deg \mu$, then $m = r + \alpha l$. By choosing a suitable basis of V , $\rho(x)$ for $x \in K'$, is a block diagonal matrix in $SO(m)$:

$$(3.8) \quad \rho(x) = \begin{pmatrix} \tau(x) & & & \\ & \mu(x) & & \\ & & \ddots & \\ & & & \mu(x) \end{pmatrix} \in SO(r) \times SO(l) \times \dots \times SO(l).$$

When μ is restricted to the subgroup $H' \subset K'$, it is not irreducible any more. Let m_0 be the multiplicity of the trivial representation in the restriction $\mu|_{H'}$. Hence for any element $y \in H'$, $\mu(y) \in SO(l - m_0) \subset SO(l)$ and $SO(l - m_0)$ is embedded as the upper left $(l - m_0) \times (l - m_0)$ block in $SO(l)$.

With the explicit description of the embeddings, we can prove the following proposition on the Weyl group of the new examples.

Proposition 3.9. *The Weyl group $\mathcal{W}$ is isomorphic to $\mathbb{Z}_2 \times \mathbb{Z}_2$.*

Proof : For any element $x \in K'$ and $A \in SO(n-m)$, let $M(x, A)$ denote the block diagonal matrix $\text{diag}(\rho(x), A)$ with $\rho(x) \in SO(m)$. If $x \in H'$, A can be considered as a matrix in $SO(n-m+1)$ since $\rho(x) \in SO(m-1)$.

First notice that w_+ can be represented by the an element $a \in K^+ = \rho(H') \times SO(n-m+1)$ which is not in H , but $a^2 \in H$. Let

$$(3.10) \quad w_+ = M(\text{id}, \begin{pmatrix} -1 & & & \\ & -1 & & \\ & & I_{n-m-1} & \end{pmatrix}) = a = \begin{pmatrix} I_{m-1} & & & \\ & -1 & & \\ & & -1 & \\ & & & I_{n-m-1} \end{pmatrix},$$

where I_k is the $k \times k$ identity matrix.

Suppose $b = M(x, A)$ is a representative of w_- , i.e., $b \in N_{K^-}(H)$ and b^2 but not $b \in H$. In the following we will determine the element b in three different cases depending on the class one representation μ .

CASE 1: μ is of real type and $m_0 = 1$. In each $W = W_i (i = 1, \dots, \alpha)$, let v be a unit vector fixed by $\mu(H')$ (or equivalently by H) and X be its orthogonal complement. Then $\dim X = l-1$. Since $b \in N_{K^-}(H)$, $b.v$ is also fixed by H and has the same length as v , i.e., $b.v = \pm v$. Since b is an orthogonal transformation, b maps X to itself, i.e., when restricted on W , $b = \begin{pmatrix} b_1 & 0 \\ 0 & \det b_1 \end{pmatrix}$ where $b_1 \in O(l-1)$. Therefore the representative b has the following matrix form:

$$(3.11) \quad w_- = b = \begin{pmatrix} A_1 & & & & & \\ & A_2 & & & & \\ & & \det A_2 & & & \\ & & & \ddots & & \\ & & & & A_2 & \\ & & & & & \det A_2 \\ & & & & & & I_{n-m} \end{pmatrix},$$

where $A_1 = \tau(x) \in SO(r)$, $A_2 = b_1 \in O(l-1)$ and there are α copies of $\begin{pmatrix} A_2 & \\ & \det A_2 \end{pmatrix}$.

CASE 2: μ is not of real type and $m_0 = 2$. In this case we have $\mu = \nu \oplus \nu^*$ where ν is a complex class one representation for the pair (K', H') . If ν is of quaternionic type, then $\nu^* = \nu$ and $\mu = \nu \oplus \nu$. As in Case 1, we have $W = X \oplus_\perp Y$ where Y is the 2 dimensional subspace fixed by $\mu(H')$ and X is its orthogonal complement. The orthogonal transformation $\mu(x)$ maps X and Y to themselves and the matrix $\mu(x)$ has the form $\begin{pmatrix} b_1 & 0 \\ 0 & b_2 \end{pmatrix} \in S(O(l-2) \times O(2))$. Since $\mu(x)^2 \in H$ implies $b_2^2 = I_2$, b_2 is a symmetric matrix. Since $\mu = \nu \oplus \nu^*$, b_2 commutes

with the matrix $\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ which implies $b_2 = \pm I_2$. Therefore b also has the matrix form as in (3.11).

CASE 3: μ is not in Case 1 or Case 2. From the classification of class one representations in Theorem A.2 and their types in Proposition A.19, the only representation of this type is $(K', H') = (Sp(k), Sp(k-1))$ where μ has the highest weight $p\varpi_1 + q\varpi_2$ with $p \geq 1$. Here ϖ_1 and ϖ_2 are the 1st and 2nd fundamental weights of $Sp(n)$. Since ρ is a faithful representation, we have $x \in K' \cap N_{K'}(H') - H'$ and $x^2 \in H'$. Take $x = -I_k \in Sp(k)$, then x satisfies these restrictions. If we view $Sp(k)$ as a subgroup of $SO(4k)$, then μ is contained in the restriction of some class one representation ν of $SO(4k)$ to $Sp(k)$ by Theorem A.1. The representation space of ν is consisted of homogeneous harmonic polynomials, so the image $\nu(x)$ is $\pm \text{Id}$ depending on the parity of the degree of the polynomials. Therefore $\mu(x)$ is equal to $\pm \text{Id}$ and the element w_- can be represented by the following matrix:

$$(3.12) \quad w_- = b = \begin{pmatrix} A_1 & & \\ & \varepsilon I_{\alpha l} & \\ & & I_{n-m} \end{pmatrix},$$

where $A_1 = \tau(x) \in SO(r)$ and $\mu(x) = \varepsilon I_l$ with $\varepsilon = \pm 1$.

In each case, from the given representatives of $w_{\pm}$, it is easy to check that $w_+w_- = w_-w_+$ which is not an element in H . Thus $\mathcal{W} = \langle w_-, w_+ \rangle \cong \mathbb{Z}_2 \times \mathbb{Z}_2$. $\square$

Remark 3.13. Let a be an element in $N(H)_0$ which does not lie in $N(K^-)$ or $N(K^+)$ and let $\bar{M}$ be the cohomogeneity one manifold defined by the group diagram $H \subset \{K^-, aK^+a^{-1}\} \subset G$. As pointed out in Remark 2.3, M and $\bar{M}$ usually have different Weyl groups and different invariant metrics though they are G -equivariantly diffeomorphic. Therefore, if one family of invariant metrics does not admit non-negative curvature, it does not necessarily follow that the other family is obstructed as well. In our example, if the multiplicity of the trivial representation in $\mu|_{H'}$ is equal to one, i.e., $m_0 = 1$, then our arguments which show the obstructions work for all equivalent diagrams. This is the case for most of the class one representations of spherical pairs, see, for example, Theorem A.2. On the other hand, if $m_0 \geq 2$, then we have to put the further restriction on the diagram: the Lie algebra of K^+ contains the subspace $\text{span}\{E_{m,m+1}, \dots, E_{m,n}\}$. Here we use $E_{i,j}$, $i \neq j$, to denote the skew-symmetric matrix having 1 in the i, j -entry, -1 in the j, i -entry and zero otherwise.

Using the explicit representatives of the generators of the Weyl group, we have the following smoothness condition of an invariant metric on M .

Lemma 3.14. *For any G -invariant metric g on M , let $h(t)$ be the length of the Killing vector field generated by $E_{m,m+1}$ along the geodesic $c(t)$. Then $h(t)$ is an even function with $h(0) \neq 0$ and $h(L) = 0$.*

Proof: The fact that $E_{m,m+1}$ lies in the Lie algebra of K^+ but not K^- implies that $h(0) \neq 0$ and $h(L) = 0$. The generator w_- is a reflection of $c(t)$ at the point p_- and maps $c(t)$ to $c(-t)$. The induced map dw_- takes $T_{c(t)}M$ to the tangent space $T_{c(-t)}M$. From the matrix form (3.11) and (3.12) of a representative of w_- , we have $dw_-(E_{m,m+1}^*(t)) = \pm E_{m,m+1}^*(-t)$. Therefore $h(t) = h(-t)$, i.e., $h(t)$ is an even function. $\square$

4. RESTRICTIONS ON THE METRIC ALONG THE NORMAL GEODESIC

In this section, we begin the study of the invariant metrics in our examples. In general, the family of G -invariant metrics on M is very large. There are many rigidity results for positively curved metrics. For example, in even dimensions, L. Verdiani classified all positively curved cohomogeneity one manifolds, see [Ve2] and [Ve3]. In odd dimensions, K.Grove, B.Wilking and W.Ziller obtained a short list of cohomogeneity one manifolds which possibly have invariant positively curved metrics in [GVWZ]. Recently K.Grove, L.Verdiani and W.Ziller have succeeded in constructing positively metric on one of them in [GVZ]. On the other hand, there are few rigidity results for non-negatively curved metrics, even in the cohomogeneity one case. Recently, B.Wilking proved some fundamental rigidity theorems for non-negatively curved manifolds in a general setting which will play an important role in our proof.

Let $c : \mathbb{R} \rightarrow M$ be a geodesic and let Λ be an $(n-1)$ -dimensional family of normal Jacobi fields. The Riccati operator $L(t)$ is the endmorphism of the normal bundle $(\dot{c}(t))^\perp$ defined by $L(t)J(t) = J'(t)$ for $J \in \Lambda$. From Theorem 9 and Corollary 10 in [Wil1], We have

Theorem 4.1 (Wilking's Rigidity Theorem). *Suppose the Riccati operator for an $(n-1)$ -dimensional family Λ of normal Jacobi fields is self-adjoint, and define the smooth subbundle Υ of $(\dot{c}(t))^\perp$ by:*

$$\Upsilon = \text{span} \{ J \in \Lambda \mid J(t) = 0 \text{ for some } t \in \mathbb{R} \}.$$

Then, if M has non-negative curvature, we have:

$$(4.2) \quad \Lambda = \Upsilon \oplus \{ J \in \Lambda \mid J \text{ is parallel} \},$$

and

$$\Upsilon(t) = \{ J(t) \mid J \in \Upsilon \} \oplus \{ J'(t) \mid J \in \Upsilon, J(t) = 0 \}.$$

A point $t_0 \in \mathbb{R}$ or $c(t_0)$ is said to be singular if $J(t_0) = 0$ for some $J \in \Upsilon$. Otherwise t_0 is said to be generic. Thus if $J \in \Lambda$ and $J(t_0) \perp \Upsilon(t_0)$ at a generic t_0 , then J is parallel along $c(t)$, $t \in \mathbb{R}$.

Remark 4.3. If there exists a subbundle $E \subset (\dot{c}(t))^\perp$ that is invariant under parallel transport, then Theorem 4.1 can also be applied to a rank E dimensional family of normal Jacobi fields in E with self-adjoint Riccati operator.

In our example, let $c(t)$ be the minimal geodesic between $B_\pm$, and $X^*(c(t))$, $X \in \mathfrak{h}^\perp$, an $(n-1)$ dimensional family of Jacobi fields. Its Riccati operator $L(t)$ is self adjoint, since it is equal to the shape operator $-\frac{1}{2}P_t^{-1}P'_t$, see (2.6). We will apply Theorem 4.1 to a subfamily of these Jacobi fields.

Let $\mathfrak{g} = \mathfrak{so}(n)$ and $\mathfrak{h}$ be the Lie algebras of $G = SO(n)$ and $H = \rho(K') \times SO(n-m)$ respectively. Choose the bi-invariant inner product $Q = -\frac{1}{2}\text{Tr}$ on $\mathfrak{g}$ for which $\{E_{i,j}\}$ is an orthonormal basis and let $\mathfrak{p}$ be the orthogonal complement of $\mathfrak{h} \subset \mathfrak{g}$. First we identify some subspaces of $\mathfrak{p}$.

Let

$$\begin{aligned}
 \mathfrak{q}_0 &= \text{span} \{E_{i,j} | 1 \leq i \leq r, m+1 \leq j \leq n\} \\
 \mathfrak{q}_1 &= \text{span} \{E_{i,j} | r+1 \leq i \leq r+l, m+1 \leq j \leq n\} \\
 &\dots \\
 \mathfrak{q}_\alpha &= \text{span} \{E_{i,j} | m+1-l \leq i \leq m, m+1 \leq j \leq n\},
 \end{aligned}
 \tag{4.4}$$

and $\mathfrak{q} = \mathfrak{q}_0 + \mathfrak{q}_1 + \dots + \mathfrak{q}_\alpha$. We write the last subspace $\mathfrak{q}_\alpha$ as a sum of two subspaces as follows:

$$\begin{aligned}
 \mathfrak{n}_1 &= \text{span} \{E_{i,j} | m+1-l \leq i \leq m-1, m+1 \leq j \leq n\}, \\
 \mathfrak{n}_2 &= \text{span} \{E_{m,j} | m+1 \leq j \leq n\}.
 \end{aligned}
 \tag{4.5}$$

Let $\mathfrak{q}^\perp$ be the Q -orthogonal complement of $\mathfrak{q}$ in $\mathfrak{p}$. Since $\mathfrak{q}^\perp$ is the fixed point set by the isotropy action of the subgroup $SO(n-m) \subset H = \rho(H') \times SO(n-m)$, Schur's lemma implies that the Killing vector field X^* , $X \in \mathfrak{q}^\perp$, is orthogonal along $c(t)$ to Y^* for any $Y \in \mathfrak{q}$.

Terminology. In the rest of the paper, for any two subspaces $\mathfrak{p}_1, \mathfrak{p}_2 \subset \mathfrak{p}$, the notation $\mathfrak{p}_1^* \perp \mathfrak{p}_2^*$ means that any Killing vector field generated by an element in $\mathfrak{p}_1$ is orthogonal to any Killing vector field generated by an element in $\mathfrak{p}_2$ along $c(t)$.

Since parallel translation commutes with the action of Ad_H , and since $\mathfrak{q}^\perp$ is the fixed point set of $SO(n-m)$ in H , it follows that $(\mathfrak{q}^\perp)^*$, and hence also $\mathfrak{q}^*$, is invariant under parallel

translation. By the same reasoning, $P^{-1}P'$ preserves $\mathfrak{q}^*$ and thus $\mathfrak{q}^*$ forms a self adjoint family of Jacobi fields to which we can thus apply Theorem 4.1.

We determine the component Υ in the splitting (4.2) of the Jacobi fields $\mathfrak{q}^*$. $p_+ = c(L)$ is a singular point. The element w_+ fixes p_+ and reflects $c(t)$ about p_+ . Let $q_- = c(2L) = w_+(p_-) \in B_-$, then the isotropy subgroup at q_- is $K_1^- = \text{Ad}_{w_+}K^-$ with Lie algebra $\mathfrak{k}_1^- = \text{Ad}_{w_+}\mathfrak{k}^-$. Similarly, w_- fixes p_- and reflects $c(t)$ about p_- . Let $q_+ = c(-L) = w_-(p_+) \in B_+$, then q_+ has isotropy subgroup $K_1^+ = \text{Ad}_{w_-}K^+$ with Lie algebra $\mathfrak{k}_1^+ = \text{Ad}_{w_-}\mathfrak{k}^+$. Since $w_-w_+ = w_+w_-$, the image of q_- under the reflection w_- about p_- is $w_-(q_-) = w_-w_+(p_-) = w_+w_-(p_-) = w_+(p_-) = q_-$, i.e., $c(2L) = c(-2L)$. Therefore $c(t)$ is a closed geodesic with period $4L$ and the singular points are $p_+ = c(L)$ and $q_+ = c(3L)$. The vanishing Killing vector fields are those generated by the vectors in the Lie algebras of the isotropy subgroups at singular points. Notice that if $X \in \mathfrak{q}$, then $X^*(p_-) \neq 0$ and $X^*(q_-) \neq 0$. Theorem 4.1 implies

Lemma 4.6. *If $Y \in \mathfrak{q}$ such that $Y^*(p_-) \perp X^*(p_-)$ for all $X \in \mathfrak{n}_2$, then Y^* is a parallel Jacobi field along $c(t)$.*

In the following, we prove some properties of the invariant metrics g on M under the non-negative sectional curvatures assumption.

First we observe

Lemma 4.7. *Suppose (M, g) is non-negatively curved, then $\mathfrak{q}_0^*$ is orthogonal to $(\mathfrak{q}_1 + \cdots + \mathfrak{q}_\alpha)^*$ along $c(t)$.*

Proof : At the generic point $p_- = c(0)$, the metric g restricted to the singular orbit $B_- \cong G/K^-$ is Ad_{K^-} -invariant. The actions of Ad_{K^-} on $\mathfrak{q}_0$ and $\mathfrak{q}_i$ ($i > 0$), are $\tau \otimes \rho_{n-m}$ and $\mu \otimes \rho_{n-m}$ respectively, where ρ_{n-m} is the standard representation of $SO(n-m)$ on $\mathbb{R}^{n-m}$. Since τ does not contain μ as a subrepresentation, $\mathfrak{q}_0^*$ is orthogonal to $\mathfrak{q}_i^*$ at p_- . In particular $\mathfrak{q}_0^*$ is orthogonal to $\mathfrak{n}_2^*$, so any Killing vector field generated by a vector in $\mathfrak{q}_0$ is parallel along $c(t)$. Hence $\mathfrak{q}_0^*$ is orthogonal to $(\mathfrak{q}_1 + \cdots + \mathfrak{q}_\alpha)^*$ along $c(t)$. $\square$

Next we study the metric g on the space $(\mathfrak{q}_1 + \cdots + \mathfrak{q}_\alpha)^*$. Recall $h^2(t) = g(E_{m,m+1}^*, E_{m,m+1}^*)$ is an even function with $h(0) \neq 0$ and W.L.O.G. we may assume that $h(0) = 1$.

4.1. μ is of real type or the multiplicity of μ in ρ is one. We first consider the case where μ is of real type. We denote $E_{r+(i-1)l+a,m+\xi}$ by $E_{a,i,\xi}$ for $a = 1, \dots, l$, $i = 1, \dots, \alpha$ and $\xi = 1, \dots, n-m$.

Since Ad_{K^-} commutes with $P(0)$, the restriction of $P(0)$ to $\mathfrak{q}_i$, composed with the projection to $\mathfrak{q}_j$, is an equivalence between the K^- irreducible representations $\mathfrak{q}_i$ and $\mathfrak{q}_j$. Since

they are orthogonal, Schur's Lemma implies that $P(0)(E_{a,i,\xi}) = f_{i,j}E_{a,j,\xi}$ for some constant $f_{i,j} \in \mathbb{R}$. Furthermore, $f_{i,j} = f_{j,i}$ from the Q -symmetry of P_t . In terms of inner product of Killing vector fields, we have

$$(4.8) \quad f_{i,j} = g(E_{1,i,1}^*, E_{1,j,1}^*)_{c(0)}.$$

The assumption $h(0) = 1$ implies that $f_{\alpha,\alpha} = 1$.

Lemma 4.9. *Suppose (M, g) is non-negatively curved, then we have*

- (1) $E_{a,i,\xi}^*$ is a parallel Jacobi field along $c(t)$ if $a \neq l$;
- (2) $E_{a,i,\xi}^*$ is orthogonal to $E_{b,j,\zeta}^*$ along $c(t)$ if $a \neq b$ or $\xi \neq \zeta$;
- (3) $E_{a,i,\xi}^*$ has the same length as $E_{a,i,\zeta}^*$ along $c(t)$;
- (4) At the point $p_- = c(0)$, $P_0(E_{a,i,\xi}) = \sum_{j=1}^{\alpha} E_{a,j,\xi} f_{i,j}$.

Proof : At the generic point $c(0)$, the Ad_K - actions on $\mathfrak{q}_i$ and $\mathfrak{q}_j$ are equivalent and given by the irreducible representation $\mu \otimes \varrho_{n-m}$, from Schur's lemma and the fact that μ is of real type or the multiplicity of μ in ρ is one, $E_{a,i,\xi}^*(0)$ is orthogonal to $\Upsilon(0) = \text{span} \{E_{l,\alpha,\zeta}^*(0) | \zeta = 1, \dots, n-m\}$ for $a \neq l$. Hence it is a parallel vector field from the last part of Theorem 4.1 which proves (1).

On each principal orbit $M_t \cong G/H$, Ad_H acts on each $\mathfrak{q}_i$ ($i > 0$), by the representation $\text{Res}_{H'}^{K'}(\mu) \otimes \varrho_{n-m}$. By Schur's lemma we have $E_{a,i,\xi}^*$ is orthogonal to $E_{b,j,\zeta}^*$ along $c(t)$ if $\xi \neq \zeta$ and $E_{a,i,\xi}^*$ has the same length as $E_{a,i,\zeta}^*$. This proves (3) and one case of (2) where $\xi \neq \zeta$.

Suppose $\zeta = \xi$ and $a \neq b$. If none of a or b is equal to l , then the two vector fields $E_{a,i,\xi}^*$ and $E_{b,j,\xi}^*$ are parallel from (1). Using Schur's lemma again and the fact that $a \neq b$, they are orthogonal to each other at $c(0)$ and then along the normal geodesic $c(t)$.

If one of a and b , say b , is equal to l , then $E_{a,i,\xi}^*$ is a parallel vector field. Write $E_{l,j,\xi}^*(0) = (E_{l,j,\xi} - \lambda E_{l,\alpha,\xi})^*(0) + \lambda E_{l,\alpha,\xi}^*(0)$, where the constant λ is determined by the following equation:

$$g(E_{l,j,\xi}^*, E_{l,\alpha,\xi}^*)_{c(0)} = \lambda g(E_{l,\alpha,\xi}^*, E_{l,\alpha,\xi}^*)_{c(0)}.$$

Thus $(E_{l,j,\xi} - \lambda E_{l,\alpha,\xi})^*(0) \perp \Upsilon(0)$ and hence $(E_{l,j,\xi} - \lambda E_{l,\alpha,\xi})^*$ is a parallel vector field. Furthermore, $E_{a,i,\xi}^*$ is orthogonal to $(E_{l,j,\xi} - \lambda E_{l,\alpha,\xi})^*$ at $c(0)$, so they are orthogonal to each other along $c(t)$. Thus $E_{a,i,\xi}^*$ is orthogonal to $E_{l,j,\xi}^*$ along $c(t)$.

The formula of $P_0(E_{a,i,\xi})$ in (4). follows easily from the defining equation (4.8) of $f_{i,j}$ and (2). $\square$

From Lemma 4.9 above, the restriction of the endomorphism P on $(\mathfrak{q}_1 + \cdots + \mathfrak{q}_\alpha)^*$ at $t = 0$ has the following matrix form:

$$(4.10) \quad P_0 = \begin{pmatrix} f_{1,1}I_l & \cdots & f_{1,\alpha}I_l \\ \vdots & \ddots & \vdots \\ f_{\alpha,1}I_l & \cdots & f_{\alpha,\alpha}I_l \end{pmatrix}.$$

We have seen that there are plenty of parallel Killing vector fields in $(\mathfrak{q}_1 + \cdots + \mathfrak{q}_\alpha)^*$. Using these parallel vector fields, we can determine the restriction of P_t on $(\mathfrak{q}_1 + \cdots + \mathfrak{q}_\alpha)^*$.

Theorem 4.11. *If the cohomogeneity one manifold (M, g) has non-negative sectional curvature and the class one representation μ is of real type, then for any $i = 1, \dots, \alpha$ and $\xi = 1, \dots, n - m$, we have*

$$(4.12) \quad \begin{aligned} (1) \quad & P_t(E_{a,i,\xi}) = \sum_{j=1}^{\alpha} f_{i,j} E_{a,j,\xi}, \text{ for } a = 1, \dots, l-1; \\ (2) \quad & P_t(E_{l,i,\xi}) = \sum_{j=1}^{\alpha} p_{i,j}(t) E_{l,j,\xi} \text{ and } p_{i,j}(t) \text{ is defined as} \\ & p_{i,j}(t) = (h^2(t) - 1)a_i a_j + f_{i,j}, \end{aligned}$$

where $a_i = f_{i,\alpha}$ and $a_\alpha = 1$.

Proof: Part (1) is obvious since every component of both $E_{a,i,\xi}^*$ and $E_{a,j,\xi}^*$ are parallel vector fields along $c(t)$ if $a \leq l-1$.

For part (2), let

$$X_i = E_{l,i,\xi} - a_i E_{l,\alpha,\xi}, \quad i = 1, \dots, \alpha.$$

Then $X_i \in \mathfrak{q}_1 + \cdots + \mathfrak{q}_\alpha$ and generates a Killing vector field X_i^* along $c(t)$. By the definition $a_i = f_{i,\alpha}$, the defining equation of $f_{i,\alpha}$ in (4.8) and (4) in Lemma 4.9, we have

$$g(X_i^*, E_{l,\alpha,\xi}^*)_{c(0)} = 0,$$

or $X_i^*(0) \perp \Upsilon(0)$. Therefore X_i^* is a parallel vector field. By the formula (2.6) of the shape operator, we have

$$P'_t(X_i) = 0 \quad \forall t \in \mathbb{R}.$$

Since $P_t(E_{l,i,\xi}) = \sum_{j=1}^{\alpha} p_{i,j}(t) E_{l,j,\xi}$ for some functions $p_{i,j}(t)$, we have

$$\begin{aligned} P'_t(E_{l,i,\xi} - a_i E_{l,\alpha,\xi}) &= P'_t(E_{l,i,\xi}) - a_i P'_t(E_{l,\alpha,\xi}) \\ &= \sum_{j=1}^{\alpha} p'_{i,j}(t) E_{l,j,\xi} - a_i \sum_{j=1}^{\alpha} p'_{\alpha,j}(t) E_{l,j,\xi} = 0. \end{aligned}$$

Therefore we have the following system of ordinary differential equations for $p_{i,j}(t)$:

$$(4.13) \quad p'_{i,j}(t) - a_i p'_{\alpha,j}(t) = 0. \quad \forall i, j = 1, \dots, \alpha.$$

One easily sees that it has the solution

$$p_{i,j}(t) = a_i a_j h^2(t) + f_{i,j} - a_i f_{j,\alpha}$$

which finishes our proof. $\square$

Now we consider the case when the multiplicity of μ in ρ is one, i.e., $\alpha = 1$. Since $P_{1,1}(0)$ is symmetric and Ad_K -equivariant, the off-diagonal terms in the formulas (2.9) and (2.10) of equivariant endomorphisms vanish, i.e., $P_{1,1}(0) = f_{1,1}I_l$ with some constant $f_{1,1} \in \mathbb{R}$. From a similar argument as in the previous case, we have

Theorem 4.14. *If the cohomogeneity one manifold (M, g) has non-negative sectional curvature and the multiplicity of the class one representation μ in ρ is one, then for any $\xi = 1, \dots, n - m$, we have*

- (1) $P_t(E_{a,1,\xi}) = f_{1,1}E_{a,1,\xi}$, if $a = 1, \dots, l - 1$;
- (2) $P_t(E_{l,1,\xi}) = p_{1,1}(t)E_{l,1,\xi}$ and $p_{1,1}(t) = (h^2(t) - 1)f_{1,1}^2 + f_{1,1}$.

4.2. μ is of complex or quaternionic type. We consider the complex case first and the quaternionic case will follow easily.

Let $l = 2p$ and $\beta = 2\alpha$. Recall that W_i is the subspace of V such that $\rho|_{W_i} = \mu$. Choose an orthonormal basis $\{e_1, \dots, e_{2p}\}$ of W_i such that $\mu(x)$ has the form $\begin{pmatrix} A & -B \\ B & A \end{pmatrix}$ with A, B being $p \times p$ matrices and $\mu(H')$ fixes the two vectors e_p and e_{2p} . Under this basis any $\text{Ad}_{\mu(K')}$ -equivariant endomorphism has the block-form $\begin{pmatrix} aI_p & -bI_p \\ bI_p & aI_p \end{pmatrix}$ as in (2.9) with constants $a, b \in \mathbb{R}$. Using the fact that the endomorphism commutes with the rotation in the $\{e_p, e_{2p}\}$ plane, we may assume that $E_{m,m} \in K^+$.

Since μ is of complex type, the Ad_K -equivariant map $P(0)$ has a block form and the (i, j) -block is given by

$$\begin{pmatrix} f_{2i-1,2j-1}I_p & f_{2i,2j-1}I_p \\ f_{2i-1,2j}I_p & f_{2i,2j}I_p \end{pmatrix},$$

where $f_{a,b} \in \mathbb{R}$ is constant for $a, b = 1, \dots, \beta = 2\alpha$ and satisfies the following identities:

(4.15)

$$f_{2i-1,2j-1} = f_{2i,2j}, \quad f_{2i,2j-1} + f_{2i-1,2j} = 0, \quad f_{2i-1,2j-1} = f_{2j-1,2i-1}, \quad f_{2i,2j-1} + f_{2j,2i-1} = 0.$$

The last two are due to the fact that $P(0)$ is Q -symmetric.

Similar to the case when μ is of real type, we define $E_{a,i,\xi} = E_{r+(i-1)p+a,m+\xi}$ for $a = 1, \dots, p$, $i = 1, \dots, \beta$ and $\xi = 1, \dots, n - m$. Then we have

Theorem 4.16. *If the cohomogeneity one manifold (M, g) has non-negative sectional curvature and the class one representation μ is of complex type, then for any $i = 1, \dots, \beta$ and $\xi = 1, \dots, n - m$, we have*

$$\begin{aligned}
(1) \quad & P_t(E_{a,i,\xi}) = \sum_{j=1}^{\beta} f_{i,j} E_{a,j,\xi}, \text{ if } a = 1, \dots, p-1; \\
(2) \quad & P_t(E_{p,i,\xi}) = \sum_{j=1}^{\beta} p_{i,j}(t) E_{p,j,\xi} \text{ and } p_{i,j}(t) \text{ is defined as} \\
(4.17) \quad & p_{i,j}(t) = (h^2(t) - 1)a_i a_j + f_{i,j}, \\
& \text{where } a_i = f_{i,\beta}.
\end{aligned}$$

Now we consider the case when μ is of quaternionic type and the multiplicity of μ in ρ is bigger than one. Let $l = 4p$ and $\beta = 4\alpha$.

From the formula (2.10) of equivariant endomorphisms in this case and a similar argument in the complex case, $P(0) = (f_{a,b} I_p)_{1 \leq a, b \leq \beta}$ where $f_{a,b} \in \mathbb{R}$ is constant and satisfies the following identities:

$$\begin{aligned}
(4.18) \quad & f_{4i-3,4j-3} = f_{4i-2,4j-2} = f_{4i-1,4j-1} = f_{4i,4j} = f_{4j,4i} \\
& -f_{4i-2,4j-3} = f_{4i-3,4j-2} = f_{4i,4j-3} = -f_{4i-1,4j} = f_{4j-1,4i} \\
& -f_{4i-1,4j-3} = -f_{4i,4j-2} = f_{4i-3,4j-1} = f_{4i-2,4j} = -f_{4j-2,4i} \\
& -f_{4i,4j-3} = f_{4i-1,4j-2} = -f_{4i-2,4j-1} = f_{4i-3,4j} = -f_{4j-3,4i}
\end{aligned}$$

for $i, j = 1, \dots, \alpha$.

We denote $E_{r+(i-1)p+a,m+\xi}$ by $E_{a,i,\xi}$ for $a = 1, \dots, p$, $i = 1, \dots, \beta$ and $\xi = 1, \dots, n-m$, and then we have

Theorem 4.19. *If the cohomogeneity one manifold (M, g) has non-negative sectional curvature and the class one representation μ is of quaternionic type, then for any $i = 1, \dots, \beta$ and $\xi = 1, \dots, n-m$, we have*

$$\begin{aligned}
(1) \quad & P_t(E_{a,i,\xi}) = \sum_{j=1}^{\beta} f_{i,j} E_{a,j,\xi}, \text{ if } a = 1, \dots, p-1; \\
(2) \quad & P_t(E_{p,i,\xi}) = \sum_{j=1}^{\beta} p_{i,j}(t) E_{p,j,\xi} \text{ and } p_{i,j}(t) \text{ is defined as} \\
(4.20) \quad & p_{i,j}(t) = (h^2(t) - 1)a_i a_j + f_{i,j}, \\
& \text{where } a_i = f_{i,\beta}.
\end{aligned}$$

5. PROOF OF THEOREM 3.1

In this section, we will develop some contradiction from the assumption that the manifold (M, g) is non-negatively curved. First we list some lemmas which will be used in the proof of Theorem 3.1.

Lemma 5.1. *The image $\mu(K') \subset SO(l)$ does not act transitively on the sphere $\mathbb{S}^{l-1} = SO(l)/SO(l-1)$.*

Proof : Recall that $l = \deg \mu \geq k + 2$. If $\mu(K')$ could act transitively on $\mathbb{S}^{l-1}$, then K' would act transitively on both $\mathbb{S}^k$ and $\mathbb{S}^{l-1}$ and $l - 1 \geq k + 1$. By the classification of the transitive action on the spheres, we have that (K', H') is either $(SO(7), SO(6))$ with $\mu(SO(7)) = Spin(7) \subset SO(8)$ or $(SO(9), SO(8))$ with $\mu(SO(9)) = Spin(9) \subset SO(16)$. But in both cases, by the classification in Theorem A.2, μ is not a class one representation for the pair (K', H') which is a contradiction. $\square$

The following was already used in [GVWZ] and we state it as a lemma without proof.

Lemma 5.2. *Suppose $f(t)$ is a C^2 non constant even function on $(-\varepsilon, \varepsilon)$ for some $\varepsilon > 0$ with $f(0) = 0$, then there is no such constant $\gamma \geq 0$ that satisfies the following inequality:*

$$(5.3) \quad \gamma^2(f(t))^2 - (f'(t))^2 \geq 0$$

We will compute the sectional curvatures for some 2-planes in our examples. The formula in terms of P_t is well established in [GZ2] and we quote it for the convenience of the reader.

Theorem 5.4 (Grove-Ziller). *If $X, Y \in \mathfrak{p}$, the sectional curvatures of M at $c(t)$ are determined by*

$$\begin{aligned} (a) \quad g(R(X, Y)X, Y) &= Q(A_-(X, Y), [X, Y]) - \frac{3}{4}Q(P[X, Y]_{\mathfrak{p}}, [X, Y]_{\mathfrak{p}}) \\ &\quad + Q(A_+(X, Y), P^{-1}A_+(X, Y)) - Q(A_+(X, X), P^{-1}A_+(Y, Y)) \\ &\quad + \frac{1}{4}Q(P'X, Y)^2 - \frac{1}{4}Q(P'X, X)Q(P'Y, Y) \\ (b) \quad g(R(X, Y)T, Y) &= -\frac{1}{2}Q(P'X, P^{-1}A_+(Y, Y)) + \frac{1}{2}Q(P'Y, P^{-1}A_+(X, Y)) \\ &\quad + \frac{3}{4}Q([X, Y], P'Y) \\ (c) \quad g(R(X, T)X, T) &= Q((-\frac{1}{2}P'' + \frac{1}{4}P'P^{-1}P')X, X). \end{aligned}$$

Remark 5.5. In the formulas above, $[X, Y]_{\mathfrak{p}}$ is the $\mathfrak{p}$ -component of $[X, Y]$ and $A_{\pm} : \mathfrak{p} \times \mathfrak{p} \longrightarrow \mathfrak{g}$ are defined as

$$A_{\pm}(X, Y) = \frac{1}{2}([X, P_t Y] \mp [P_t X, Y]).$$

From Lemma 3.14, $h(t)$ is an even function with $h(0) \neq 0$ and $h(L) = 0$, so $f(t) = h^2(0) - h^2(t) = 1 - h^2(t)$ satisfies the assumptions in Lemma 5.2. We will show the inequality (5.3) holds for some constant $\gamma \geq 0$ using nonnegativity of the sectional curvature of a carefully chosen 2-plane. Theorem 3.1 then follows from Lemma 5.2.

Proof of Theorem 3.1: In the following argument, only the entries in the lower right $(n-r) \times (n-r)$ -block of $\mathfrak{g} = \mathfrak{so}(n)$ are involved, so without loss of generality we may assume that $r = 0$ or equivalently the representation ρ is a sum of α copies of μ .

CASE 1: The representation μ is of real type. Since P_0 defined in (4.10) is symmetric and positive definite, we can write

$$\begin{pmatrix} f_{1,1} & \cdots & f_{1,\alpha} \\ \vdots & \ddots & \vdots \\ f_{\alpha,1} & \cdots & f_{\alpha,\alpha} \end{pmatrix} = ADA^\top,$$

where $A = (A_{i,j})_{\alpha \times \alpha}$ is an orthogonal matrix and $D = \text{diag}(d_1, \dots, d_\alpha)$ is a diagonal matrix with positive entries. Define the following vectors in $\mathfrak{q}_1 + \cdots + \mathfrak{q}_\alpha$:

$$(5.6) \quad X^u = \sum_{i=1}^{l-1} b_i E_{i,u,1} + E_{l,u,2}$$

$$(5.7) \quad Y^u = \sum_{i=1}^{l-1} b_i E_{i,u,2} + E_{l,u,1},$$

where $u = 1, \dots, \alpha$ and $\sum_{i=1}^{l-1} b_i^2 = 1$. Further conditions of the b_i 's will be determined later on.

In the matrix A , there is a column, say the i_0 -th column, with $A_{\alpha, i_0} \neq 0$. We denote A_{u, i_0} by A_u , $u = 1, \dots, \alpha$, and define the following two vectors X, Y in $\mathfrak{p}$:

$$(5.8) \quad X = \sum_{u=1}^{\alpha} A_u X^u, \quad Y = \sum_{u=1}^{\alpha} A_u Y^u.$$

From the definitions of X^u and Y^v , it is easy to see that $[X^u, Y^u] = 0$, and if $u \neq v$, then

$$[X^u, Y^v] = \sum_{i=1}^{l-1} b_i (E_{vl, i+(u-1)l} + E_{i+(v-1)l, ul}).$$

Hence

$$\begin{aligned} [X, Y] &= \sum_{u,v=1}^{\alpha} A_u A_v [X^u, Y^v] = \sum_{i=1}^{l-1} \sum_{u \neq v} A_u A_v b_i (E_{vl, i+(u-1)l} + E_{i+(v-1)l, ul}) \\ &= 0. \end{aligned}$$

The fact that X commutes with Y makes the computation of the sectional curvature of the 2-plane spanned by X^* and Y^* easier since the first two terms in the curvature formula

(a) in Theorem 5.4 vanish. The other four terms are computed in Proposition 5.9 below. If we plug in the result of each term into the formula of the sectional curvature, we have

$$\begin{aligned} \|X^* \wedge Y^*\|^2 \sec(X^*, Y^*)_{c(t)} &= (h^2(t) - 1)^2 Q(X_0, P_t^{-1}(X_0)) - (d_{i_0} A_\alpha)^4 h^2(t) (h'(t))^2 \\ &\quad - Q(A_+(X, X), P_t^{-1} A_+(X, X)). \end{aligned}$$

Here $X_0 \in \mathfrak{p}$ is specified in Proposition 5.9 and is orthogonal to $\mathfrak{k}^-$ with respect to Q .

Since P_t is positive definite as well as P_t^{-1} , we have $Q(A_+(X, X), P_t^{-1} A_+(X, X)) \geq 0$. Therefore $\sec(X^*, Y^*) \geq 0$ implies that

$$(h^2(t) - 1)^2 Q(X_0, P_t^{-1}(X_0)) - (d_{i_0} A_\alpha)^4 h^2(t) (h'(t))^2 \geq 0.$$

The existence of the constant $\gamma \geq 0$ follows from the facts that $A_\alpha \neq 0$ and $Q(X_0, P_t^{-1}(X_0))$ is bounded from above near $t = 0$.

CASE 2: The representation μ is not of real type and the multiplicity of μ in ρ equals to 1. In this case P_0 is a scalar matrix and P_t is a diagonal matrix. It is easy to see that the proof in the previous case works if we choose $X = X^1$ and $Y = Y^1$ in (5.6) and (5.7).

CASE 3: The representation μ is not of real type and the multiplicity of μ in ρ is bigger than 1. Let $p = \frac{1}{2}l$ and $\beta = 2\alpha$ if μ is of complex type and let $p = \frac{1}{4}l$ and $\beta = 4\alpha$ if it is of quaternionic type. In both cases, we have $p \geq k + 2$. In each case we define the vector $E_{a,i,\xi}$ for $a = 1, \dots, p$, $i = 1, \dots, \beta$ and $\xi = 1, \dots, n - m$. Then similarly we can define vectors X^u and Y^u for $u = 1, \dots, \beta$ and use them to define the vectors X and Y as in CASE 1. The formulas of $P_t(X)$ and $P_t(Y)$ are obtained by Theorem 4.16 and Theorem 4.19 respectively and the rest of the proof will follow CASE 1. Note that the number of the constants b_i 's in X or Y is equal to $p - 1$. The existence of the vector X_0 which is orthogonal to $\mathfrak{k}^-$ follows from the fact that $p \geq k + 2$. $\square$

In the case where μ is real type, the non-vanishing terms in the sectional curvature of the 2-plane spanned by X and Y are computed in the following

Proposition 5.9. *For the vectors X and Y defined in (5.8), by choosing a proper value for the constant b_i , we have*

- (1) *There exists some $X_0 \in \mathfrak{p}$ which is orthogonal to $\mathfrak{k}^-$ with respect to Q such that $A_+(X, Y) = (h^2(t) - 1)X_0$;*
- (2) $A_+(X, X) = A_+(Y, Y)$;
- (3) $Q(P'_t(X), Y) = 0$;
- (4) $-\frac{1}{4}Q(P'_t(X), X)Q(P'_t(Y), Y) = -(d_{i_0} A_\alpha)^4 h^2(t) (h'(t))^2$.

Proof : First we compute the endomorphism P_t on X and Y . From the defining equations (5.6) of X^u and (5.7) of Y^u , we have

$$(5.10) \quad P_t(X^u) = \sum_{i=1}^{l-1} \sum_{r=1}^{\alpha} b_i f_{u,r} E_{i,r,1} + \sum_{r=1}^{\alpha} p_{u,r}(t) E_{l,r,2}$$

and

$$(5.11) \quad P_t(Y^v) = \sum_{j=1}^{l-1} \sum_{s=1}^{\alpha} b_j f_{v,s} E_{j,s,2} + \sum_{s=1}^{\alpha} p_{v,s}(t) E_{l,s,1},$$

then

$$\begin{aligned} [X^u, P_t(Y^v)] &= \left[\sum_{i=1}^{l-1} b_i E_{i+(u-1)l, \alpha l+1} + E_{ul, \alpha l+2}, \sum_{j=1}^{l-1} \sum_{s=1}^{\alpha} b_j f_{v,s} E_{j+(s-1)l, \alpha l+2} + \sum_{s=1}^{\alpha} p_{v,s}(t) E_{sl, \alpha l+1} \right] \\ &= \left(\sum_{i=1}^{l-1} b_i^2 f_{v,u} - p_{v,u}(t) \right) E_{\alpha l+2, \alpha l+1} + \sum_{i=1}^{l-1} \sum_{s=1}^{\alpha} b_i (f_{v,s} E_{i+(s-1)l, ul} - p_{v,s}(t) E_{i+(u-1)l, sl}), \end{aligned}$$

and

$$\begin{aligned} [P_t(X^u), Y^v] &= \left[\sum_{i=1}^{l-1} \sum_{r=1}^{\alpha} b_i f_{u,r} E_{i+(r-1)l, \alpha l+1} + \sum_{r=1}^{\alpha} p_{u,r}(t) E_{rl, \alpha l+2}, \sum_{j=1}^{l-1} b_j E_{j+(v-1)l, \alpha l+2} + E_{vl, \alpha l+1} \right] \\ &= \left(\sum_{i=1}^{l-1} b_i^2 f_{u,v} - p_{u,v}(t) \right) E_{\alpha l+2, \alpha l+1} + \sum_{i=1}^{l-1} \sum_{r=1}^{\alpha} b_i (f_{u,r} E_{vl, i+(r-1)l} - p_{u,r}(t) E_{rl, i+(v-1)l}). \end{aligned}$$

Therefore

$$(5.12) \quad A_+(X^u, Y^v) = \frac{1}{2} \sum_{i=1}^{l-1} \sum_{s=1}^{\alpha} b_i (f_{v,s} E_{i+(s-1)l, ul} + f_{u,s} E_{i+(s-1)l, vl} - p_{v,s}(t) E_{i+(u-1)l, sl} - p_{u,s}(t) E_{i+(v-1)l, sl}).$$

From the above equation, only the terms as $E_{j+(r-1)l, wl}$ have nonzero coefficients in $A_+(X, Y)$ and it is denoted by $c_{j,r,w}$. From the formula (5.12) and the bi-linearity of A_+ , we have

$$\begin{aligned} c_{j,r,w} &= \frac{1}{2} b_j \left(\sum_{v=1}^{\alpha} (f_{v,r} A_w A_v - p_{v,w}(t) A_r A_v) + \sum_{u=1}^{\alpha} (f_{u,r} A_u A_w - p_{u,w}(t) A_u A_r) \right) \\ (5.13) \quad &= b_j \left(A_w \sum_{v=1}^{\alpha} f_{v,r} A_v - A_r \sum_{v=1}^{\alpha} p_{v,w}(t) A_v \right). \end{aligned}$$

We can compute the terms in (5.13) explicitly as follows,

$$(5.14) \quad \sum_{v=1}^{\alpha} f_{v,r} A_v = \sum_{v=1}^{\alpha} \sum_{i=1}^{\alpha} A_{v,i} d_i A_{r,i} A_{v,i_0} = (A^{\tau} A D A^{\tau})_{i_0, r} = d_{i_0} A_{r, i_0} = d_{i_0} A_r$$

and

$$\begin{aligned}
\sum_{v=1}^{\alpha} p_{v,w}(t) A_v &= \sum_{v=1}^{\alpha} (a_v a_w h^2(t) A_v + f_{v,w} A_v - a_v f_{w,\alpha} A_v) \\
&= d_{i_0} A_w + (a_w h^2(t) - f_{w,\alpha}) \sum_{v=1}^{\alpha} a_v A_v \\
&= d_{i_0} A_w + a_w (h^2(t) - 1) \sum_{v=1}^{\alpha} f_{v,\alpha} A_v \\
(5.15) \quad &= d_{i_0} A_w + d_{i_0} a_w A_{\alpha} (h^2(t) - 1),
\end{aligned}$$

where the first equality follows Theorem 4.11.

By substituting the new expressions (5.14) and (5.15) back in the expression (5.13) of $c_{j,r,w}$, we have

$$(5.16) \quad c_{j,r,w} = -d_{i_0} A_r A_{\alpha} a_w b_j (h^2(t) - 1).$$

If $r \neq w$, then $E_{j+(r-1)l, wl}$ is orthogonal to $\mathfrak{k}^-$ with respect to Q . If $r = w$, by the formula (5.16) for $c_{j,r,w}$, $c_{j,r}$ is a multiple of b_j . From the assumption that the representation ρ is the direct sum of μ and the embedding in (3.8), the image $\mu_*(v)$ (for any $v \in \mathfrak{k}'$) is block-wise diagonally embedded in $\mathfrak{so}(m)$. Hence if for some r , the vector $v_r = \sum_{j=1}^{l-1} c_{j,r} E_{j+(r-1)l, rl}$ is orthogonal to $\mathfrak{k}^-$, then all vectors v_q 's are orthogonal to $\mathfrak{k}^-$. By Lemma 5.1 of the non-transitivity of the action $\mu(K')$ on the sphere $SO(l)/SO(l-1)$ and by choosing the proper values of b_i 's, v_r is orthogonal to $\mathfrak{k}^-$. Therefore $A_+(X, Y) = (h^2(t) - 1)X_0$ for some $X_0 \in \mathfrak{p}$ which is orthogonal to $\mathfrak{k}^-$ with respect to Q and (1) is proved.

By (5.6) and (5.10), we have

$$\begin{aligned}
[X^u, P_t(X^v)] &= \left[\sum_{i=1}^{l-1} b_i E_{i+(u-1)l, \alpha l+1} + E_{ul, \alpha l+2}, \sum_{j=1}^{l-1} \sum_{s=1}^{\alpha} b_j f_{v,s} E_{j+(s-1)l, \alpha l+1} + \sum_{s=1}^{\alpha} p_{v,s}(t) E_{sl, \alpha l+2} \right] \\
&= \sum_{i \neq j, \text{ OR } u \neq s} b_i b_j f_{v,s} E_{j+(s-1)l, i+(u-1)l} + \sum_{s \neq u} p_{v,s}(t) E_{sl, ul}.
\end{aligned}$$

By (5.7) and (5.11), we have the same result for $[Y^u, P_t(Y^v)]$, therefore $A_+(X, X) = [X, P_t(X)] = [Y, P_t(Y)] = A_+(Y, Y)$ which proves the formula in (2).

Next we will prove the formulas in (3) and (4) which will finish the proof of the proposition.

By (5.10) and the differential equation (4.13) for p_{ij} we have

$$P'_t(X^u) = \sum_{r=1}^{\alpha} p'_{r,u}(t) E_{l,r,2} = 2h(t)h'(t) \sum_{r=1}^{\alpha} a_u a_r E_{l,r,2},$$

so

$$\begin{aligned} Q(P'_t(X^u), Y^v) &= 2a_u h(t) h'(t) \sum_{r=1}^{\alpha} a_r \left(\sum_{j=1}^{l-1} b_j Q(E_{l,r,2}, E_{j,v,2}) + Q(E_{l,r,2}, E_{l,v,1}) \right) \\ &= 0, \end{aligned}$$

and then $Q(P'_t(X), Y) = 0$.

By taking the inner product with X^v instead of Y^v , we have

$$\begin{aligned} Q(P'_t(X^u), X^v) &= 2a_u h(t) h'(t) \sum_{r=1}^{\alpha} a_r \left(\sum_{j=1}^{l-1} b_j Q(E_{l,r,2}, E_{j,v,1}) + Q(E_{l,r,2}, E_{l,v,2}) \right) \\ &= 2a_u a_v h(t) h'(t). \end{aligned}$$

Therefore

$$\begin{aligned} Q(P'_t(X), X) &= 2 \left(\sum_{u,v=1}^{\alpha} A_u A_v a_u a_v \right) h(t) h'(t) = 2 \left(\sum_{u=1}^{\alpha} A_u a_u \right)^2 h(t) h'(t) \\ &= 2 (d_{i_0} A_{\alpha})^2 h(t) h'(t), \end{aligned}$$

where the last equality follows either from (5.14) or (5.15). Similarly for $Q(P'_t(Y), Y)$ and hence (4) follows, which finishes the proof. $\square$

Remark 5.17. In the introduction, we pointed out that there are unknown but interesting cases when $m \geq k + 2$ and $n = m + 1$. The minimal dimension of these manifolds is 15 when $k = 2$, $m = 5$ and $n = 6$. This manifold has cohomology ring different from the two 15 dimensional symmetric spaces $\mathbb{S}^{15}$ and $SO(8)/(SO(5) \times SO(3))$. The geometry of these examples will be studied in another paper.

6. COHOMOGENEITY ONE MANIFOLDS FOR $G = U(n)$ AND $Sp(n)$

In this section, we will generalize our examples to the cases where $G = U(n)$ and $Sp(n)$. First let us state the theorem in each case.

Theorem 6.1. *Let $K'/H' = \mathbb{S}^k$ with $k \geq 2$ and $\rho : K' \longrightarrow U(m)$ be a faithful representation. Suppose ρ contains a class one representation $\mu : K' \longrightarrow U(l)$ of the pair (K', H') with $2l \geq k + 2$ and the multiplicity of μ in ρ is 1. For any integer $n \geq m + 2$, set $G = U(n)$ and*

$$\begin{aligned} K^- &= \rho(K') \times U(n - m) \subset U(m) \times U(n - m) \subset U(n) \\ (6.2) \quad K^+ &= \rho(H') \times U(n - m + 1) \subset U(m - 1) \times U(n - m + 1) \subset U(n) \\ H &= \rho(H') \times U(n - m) \subset U(n), \end{aligned}$$

then the cohomogeneity one manifold M defined by the groups $H \subset \{K^-, K^+\} \subset G$ does not admit a G invariant metric with non-negative sectional curvature.

Remark 6.3. Proposition A.22 lists the complex class one representations which have dimension smaller than $\frac{1}{2}(k+2)$. It shows that if $\text{mul}(\mu, \rho) = 1$, then only the defining representations of $SU(l)$, $U(l)$, $Sp(l)$ and $Sp(l) \times U(1)$ are excluded by the above theorem. The cohomogeneity one manifolds from these representations are equivariantly diffeomorphic to the homogeneous spaces $U(n+1)/\Phi(K') \times U(n-l+1)$ where $\Phi : K' \rightarrow U(l)$ is the defining representation, so they carry non-negatively curved metrics.

Similar to Lemma 5.1 in the orthogonal case, we have the following lemma:

Lemma 6.4. *Assume that K' , H' and μ as in Theorem 6.1 with $2l \geq k+2$, then $\mu(K')$ does not act transitively on $\mathbb{CP}^{l-1} = U(l)/(U(l-1) \times U(1))$.*

Proof : From the classification of the transitive actions on complex projective spaces, see [Be2], p.195, we only need to check the pair $(SU(2), U(1))$ for $\mathbb{CP}^1$, $(U(n), U(n-1))$ for $\mathbb{CP}^{n-1}$ and $(Sp(n), Sp(n-1))$ for $\mathbb{CP}^{2n-1}$. In the first case, the subgroup $U(1) \subset SU(2)$ does not fix any vector in $\mathbb{C}^2$. In the last two cases, we have $2l = k+1$ which contradicts the assumption on l . Therefore the action of $\mu(K')$ on $\mathbb{CP}^{l-1}$ is non-transitive. $\square$

Sketch of the Proof of Theorem 6.1: Suppose ρ has the decomposition $\tau \oplus \mu$ and μ is not equivalent to any of subrepresentations in τ . Let $c(t)$ be the normal geodesic connecting the two singular orbits $B_{\pm} = G/K^{\pm}$ with $c(0) = p_- \in B_-$ and $c(L) = p_+ \in B_+$. Similar to the orthogonal case, the Weyl group is isomorphic to $\mathbb{Z}_2 \times \mathbb{Z}_2$ and the generators have the following representatives:

$$w_+ = \begin{pmatrix} I_{m-1} & & \\ & -1 & \\ & & I_{n-m} \end{pmatrix},$$

and

$$w_- = \begin{pmatrix} A_1 & & & \\ & A_2 & & \\ & & \varepsilon & \\ & & & I_{n-m} \end{pmatrix} \text{ or } \begin{pmatrix} A_1 & & & \\ & \varepsilon I_l & & \\ & & & I_{n-m} \end{pmatrix},$$

where $A_1 \in U(r)$, $A_2 \in U(l-1)$ and $\varepsilon = \pm 1$.

In addition to the matrices $\{E_{i,j}\}_{1 \leq i \neq j \leq n}$, let $F_{i,j}$ be the symmetric matrix with $i = \sqrt{-1}$ in the i, j - and j, i -entries if $i \neq j$ and $\sqrt{2}i$ in the i, i -entry if $i = j$. Then $\{E_{i,j}\}$ and $\{F_{i,j}\}$ form an orthonormal basis of the Lie algebra $\mathfrak{u}(n)$ of $U(n)$ with $Q = -\frac{1}{2}\Re \text{Tr}$, where $\Re$ takes the real part of a complex number. Without loss of generality, we may assume that

$r = 0$, i.e., $\rho = \mu$ and then $m = l$. Let $\mathfrak{p}$ be the orthogonal complement of the Lie algebra $\mathfrak{h}$ of H in the Lie algebra $\mathfrak{g} = \mathfrak{u}(n)$ of G and $\mathfrak{q}$ be the subspace of $\mathfrak{p}$ spanned by the vectors $\{E_{a,i}\}$ and $\{F_{a,i}\}$ for $a = 1, \dots, m$ and $i = m+1, \dots, n$.

Let $h^2(t) = g(E_{m,m+1}^*, E_{m,m+1}^*)_{c(t)}$ and then we may assume that $h(0) = 1$. From Schur's Lemma and Wilking's Rigidity Theorem, we have

Proposition 6.5. *Suppose that the metric g is non-negatively curved, then we have*

- (1) $P(E_{a,i}) = E_{a,i}$ and $P(F_{a,i}) = F_{a,i}$, if $a = 1, \dots, m-1$;
- (2) $P(E_{m,i}) = h^2(t)E_{m,i}$ and $P(F_{m,i}) = h^2(t)F_{m,i}$,

where $i = m+1, \dots, n$.

From the collapsing of the Killing vector field $E_{m,m+1}^*$ at p_+ and Weyl symmetry at p_- , $h(t)$ is an even function and $h(L) = 0$. Let

$$\begin{aligned} X &= \sum_{i=1}^{m-1} b_i E_{i,m+1} + E_{m,m+2} + \sum_{i=1}^{m-1} c_i F_{i,m+1} + F_{m,m+2} \\ Y &= \sum_{j=1}^{m-1} b_j E_{j,m+2} + E_{m,m+1} + \sum_{j=1}^{m-1} c_j F_{j,m+2} + F_{m,m+1}, \end{aligned}$$

where $\sum_{i=1}^{m-1} (b_i^2 + c_i^2) = 2$.

A computation shows that $[X, Y] = 0$ and results in the following claim:

Claim. *For properly chosen values of the constants b_i and c_i , we have*

- (1) *There exists some $X_0 \in \mathfrak{p}$ which is orthogonal to $\mathfrak{k}^-$, with respect to Q such that $A_+(X, Y) = (h^2(t) - 1)X_0$;*
- (2) $A_+(X, X) = A_+(Y, Y)$;
- (3) $Q(P'_t(X), Y) = 0$;
- (4) $-\frac{1}{4}Q(P'_t(X), X)Q(P'_t(Y), Y) = -4(h(t)h'(t))^2$.

The existence of X_0 follows from the non-transitivity of the $\mu(K')$ action on $\mathbb{CP}^{l-1} = U(l)/(U(l-1) \times U(1))$ proved in Lemma 6.4. The same argument as in the orthogonal case shows that the non-negativity of the sectional curvature of the 2-plane spanned by X^* and Y^* gives the desired contradiction. This completes the proof in the unitary case. $\square$

Finally we discuss the case where $G = Sp(n)$.

Theorem 6.6. *Let $K'/H' = \mathbb{S}^k$ with $k \geq 2$ and $\rho : K' \longrightarrow Sp(m)$ be a faithful representation. Suppose ρ contains a class one representation $\mu : K' \longrightarrow Sp(l)$ of the pair (K', H')*

with $4l \geq k+2$ and the multiplicity of μ in ρ is 1. For any integer $n \geq m+2$, set $G = Sp(n)$ and

$$\begin{aligned} K^- &= \rho(K') \times Sp(n-m) \subset Sp(m) \times Sp(n-m) \subset Sp(n) \\ (6.7) \quad K^+ &= \rho(H') \times Sp(n-m+1) \subset Sp(m-1) \times Sp(n-m+1) \subset Sp(n) \\ H &= \rho(H') \times Sp(n-m) \subset Sp(n), \end{aligned}$$

then the cohomogeneity one manifold M defined by the groups $H \subset \{K^-, K^+\} \subset G$ does not admit a G invariant metric with non-negative sectional curvature.

Remark 6.8. Proposition A.23 lists the quaternionic class one representations which have dimension smaller than $\frac{1}{4}(k+2)$. It show that only the standard representation of $Sp(l)$ for the pair $(Sp(l), Sp(l-1))$ is excluded by the above theorem. The cohomogeneity one manifold from this representation has non-negatively curved metric since it is equivariantly diffeomorphic to the homogeneous space $Sp(n+1)/Sp(l) \times Sp(n-l+1)$.

We have the following result on quaternionic projective spaces which is analogues to Lemma 5.1 and Lemma 6.4:

Lemma 6.9. *Assume that K' , H' and μ are as in Theorem 6.6 with $4l \geq k+2$, then $\mu(K')$ does not act transitively on $\mathbb{HP}^{l-1} = Sp(l)/(Sp(l-1) \times Sp(1))$.*

Proof : From the classification of the transitive actions on $\mathbb{HP}^{l-1}$, see [Be2], p.195, we have $\mu(K') = Sp(l)$ and then $H' = Sp(l-1)$, μ is the standard representation of $K' = Sp(l)$. However in this case, $k = 4l - 1$ which contradicts the assumption $4l \geq k+2$. $\square$

Sketch of the Proof of Theorem 6.6: The proof follows as in the complex case where $G = U(n)$ step by step. Suppose $\{1, \iota, j, \kappa\}$ is the basis of $\mathbb{H}$ over the reals such that $\iota^2 = j^2 = \kappa^2 = -1$ and $\kappa = \iota j$. Let $G_{i,j}$ denote the symmetric matrix with 1 in the i, j - and j, i -entries if $i \neq j$, and $\sqrt{2}$ in the i, i -entries. Thus $\{E_{i,j}, \iota G_{i,j}, j G_{i,j}, \kappa G_{i,j}\}$ forms an orthonormal basis of the Lie algebra $\mathfrak{sp}(n)$ of $Sp(n)$ with $Q = -\frac{1}{2}\Re\text{Tr}$. Without loss of generality, we may assume that $r = 0$, i.e., $m = l$.

Let $h^2(t) = g(E_{m,m+1}^*, E_{m,m+1}^*)_{c(t)}$ and we may assume that $h(0) = 1$. For the endomorphism P_t , one proves the following proposition which is similar to the orthogonal and complex cases.

Proposition 6.10. *Suppose that the metric g is non-negatively curved, then we have*

- (1) $P(E_{a,i}) = E_{a,i}$ and $P(\theta G_{a,i}) = \theta G_{a,i}$, if $a = 1, \dots, m-1$;
- (2) $P(E_{m,i}) = h^2(t)E_{m,i}$ and $P(\theta G_{m,i}) = h^2(t)\theta G_{m,i}$,

where $i = m+1, \dots, n$ and θ can be one of ι, j and κ .

Furthermore, $h(t)$ is an even function with $h(L) = 0$. To get the desired contradiction we choose the vectors X and Y as follows:

$$\begin{aligned} X &= \sum_{i=1}^{m-1} b_i E_{i,m+1} + E_{m,m+2} + \sum_{i=1}^{m-1} c_i G_{i,m+1} + c G_{m,m+2}, \\ Y &= \sum_{j=1}^{m-1} b_j E_{j,m+2} + E_{m,m+1} + \sum_{j=1}^{m-1} c_j G_{j,m+2} + c G_{m,m+1}, \end{aligned}$$

where $c = \iota + j + \kappa$, b_i 's are real numbers and c_i 's are pure quaternionic numbers, i.e. the real part is zero. These constants satisfy the equation $1 - c^2 - \sum_{i=1}^{l-1} (b_i^2 - c_i^2) = 0$.

A computation shows that $[X, Y] = 0$ and the following claim:

Claim. *For properly chosen values of the constants b_i and c_i , we have*

- (1) *There exists some $X_0 \in \mathfrak{p}$ which orthogonal to $\mathfrak{k}^-$, with respect to Q such that $A_+(X, Y) = (h^2(t) - h^2(0))X_0$;*
- (2) $A_+(X, X) = A_+(Y, Y)$;
- (3) $Q(P'_t(X), Y) = 0$;
- (4) $-\frac{1}{4}Q(P'_t(X), X)Q(P'_t(Y), Y) = -16(h(t)h'(t))^2$.

The existence of X_0 follows from the non-transitivity of the $\mu(K')$ action on $\mathbb{H}P^{l-1}$ proved in Lemma 6.9. The contradiction now follows as in the orthogonal case. $\square$

APPENDIX A. CLASS ONE REPRESENTATIONS OF SPHERE GROUP PAIRS

In this appendix, we shall classify all class one representations of spherical pairs. Except for the last section, all representations are considered over the complex numbers.

First we reduce our classification to the case where the group action on the sphere is almost effective. Suppose K, H are compact Lie groups, K is connected and $K/H = \mathbb{S}^{n-1}$ with $n \geq 2$. If the K action is not almost effective, then let C be the ineffective kernel, i.e. the maximal normal subgroup of K contained in H . Thus we can write $K = C \times K_1$ and $H = C \times H_1$. Suppose $\mu \otimes \tau$ is a class one representation of the pair (K, H) , i.e., μ and τ are irreducible representations of C and K_1 respectively and $\text{Res}(\mu \otimes \tau)$ fixes a non-zero vector, i.e., the trivial representation of $C \times H_1$ appears in the decomposition of $\mu \otimes \tau$. Therefore μ is the trivial representation of C and τ is a class one representation of the pair (K_1, H_1) . Therefore in the rest, we only consider almost effective action on the spheres.

The classification of transitive and effective action on spheres by connected compact Lie groups is well known, see for example Page 179 in [Be1] or Page 195 in [Be2]. Using representation theory of compact Lie groups, it is easy to extend the classification to the almost effective case:

- $SO(n)/SO(n-1) = \mathbb{S}^{n-1} (n \geq 3)$,
- $SU(n)/SU(n-1) = \mathbb{S}^{2n-1} (n \geq 2)$,
- $U(n)/U(n-1)_m = \mathbb{S}^{2n-1} (n \geq 1 \text{ and } m \in \mathbb{Z})$,
- $Sp(n)/Sp(n-1) = \mathbb{S}^{4n-1} (n \geq 1)$,
- $Sp(n) \times Sp(1)/(Sp(n-1) \times Sp(1)) = \mathbb{S}^{4n-1} (n \geq 1)$,
- $Sp(n) \times U(1)/(Sp(n-1) \times U(1))_m = \mathbb{S}^{4n-1} (n \geq 1 \text{ and } m \in \mathbb{Z}, m \neq 0)$,
- $G_2/SU(3) = \mathbb{S}^6$, $Spin(7)/G_2 = \mathbb{S}^7$, $Spin(9)/Spin(7) = \mathbb{S}^{15}$.

The group $U(n)$ can act on the sphere $\mathbb{S}^{2n-1}$ in different ways. For each $m \in \mathbb{Z}$, $A \in U(n)$ can act on $\mathbb{C}^n$ via $z \mapsto (\det A)^m A.z$. It induces a transitive action on $\mathbb{S}^{2n-1} \subset \mathbb{C}^n$ and the isotropy subgroup at $z = (1, 0, \dots, 0)^\top$ is

$$U(n-1)_m = \{ \text{diag}(a, B) \in U(n) \mid a^{m+1} = (\det B)^{-m}, a \in U(1) \text{ and } B \in U(n-1) \}.$$

If $n = 1$, then $U(1)$ acts on the circle $\mathbb{S}^1$ and the isotropy subgroup is $\mathbb{Z}_{m+1} (m \neq -1)$ and it gives us all almost effective actions on the circle. In Table 1, we only list the case when the $U(1)$ action on $\mathbb{S}^1$ is effective and the corresponding results in the almost effective case easily follows.

Similarly the group $Sp(n) \times U(1)$ has different transitive actions on $\mathbb{S}^{4n-1}$. For each $m \in \mathbb{Z}$, $(A, z) \in Sp(n) \times U(1)$ acts on $\mathbb{H}^n$ via $q \mapsto A.qz^m$. The isotropy subgroup at $q = (0, \dots, 0, 1)^\top$ is

$$Sp(n-1) \times U(1)_m = \{ (\text{diag}(A, z^{-m}), z) \in Sp(n) \times U(1) \mid z \in U(1) \text{ and } A \in Sp(n-1) \}.$$

For each spherical pair (K, H) with $K/H = \mathbb{S}^{n-1}$, the defining representation $\Phi : K \longrightarrow SO(n)$ is of class one. If the pair is $(SO(n), SO(n-1))$, then the class one representations are well known. They consist of the irreducible representations on the space of homogeneous harmonic polynomials. In fact the class one representations of the spherical pairs are closely related to these representations as stated in the following theorem.

Theorem A.1. *The representation μ is a class one representation of the spherical pair (K, H) if and only if μ is in the decomposition of $\text{Res}_K^{SO(n)} \rho$, where ρ is a class one representation of the pair $(SO(n), SO(n-1))$ and K is viewed as a subgroup of $SO(n)$ via the representation Φ .*

The proof of the above theorem is given in Section A.1.

For a compact Lie group, the irreducible representations are highest weight representations and each highest weight is a linear combination of the fundamental weights with nonnegative integers as coefficients. We list the fundamental weights for classical groups as follows.

$$\begin{aligned}
SO(n) : \quad & \varpi_1 = e_1, \dots, \varpi_{k-1} = e_1 + e_2 + \dots + e_{k-1}, \varpi_k = \frac{1}{2}(e_1 + \dots + e_k), \quad n = 2k+1, \\
& \varpi_1 = e_1, \dots, \varpi_{k-1} = \frac{1}{2}(e_1 + \dots + e_{k-1} - e_k), \varpi_k = \frac{1}{2}(e_1 + \dots + e_k), \quad n = 2k, \\
SU(n) : \quad & \varpi_1 = e_1, \dots, \varpi_{n-1} = e_1 + \dots + e_{n-1}, \text{ with } e_1 + \dots + e_n = 0, \\
Sp(n) : \quad & \varpi_1 = e_1, \dots, \varpi_n = e_1 + \dots + e_n.
\end{aligned}$$

The exceptional Lie group G_2 has two fundamental weights: ϖ_1 which is the highest weight of the 7 dimensional representation and ϖ_2 which is 14 dimensional.

The group $U(n)$ has a finite cover $SU(n) \times U(1)$ and hence its irreducible representations can be written as $\mu \otimes \phi^k$ where μ is an irreducible representation of $SU(n)$ and $\phi^k (k \in \mathbb{Z})$ is the 1 dimensional representation of $U(1)$: $v \mapsto z^k v$ for any $z \in U(1)$. Hence an irreducible representation ρ of $U(n)$ with highest weight $a_1 e_1 + \dots + a_n e_n$ ($a_1 \geq \dots \geq a_n$) is the tensor product of an irreducible representation μ of $SU(n)$ and ϕ^k of $U(1)$ where μ has highest weight $(a_1 - a_n)e_1 + \dots + (a_{n-1} - a_n)e_{n-1}$ and $k = -(a_1 + \dots + a_n)$. Note that the standard representation of $U(n)$ on $\mathbb{C}^n$, i.e., matrix multiplication from the left, has the highest weight $-e_n$.

In order to determine the class one representations for each pair (K, H) , we use the branching rules, i.e. the rules that show how an irreducible representation of K decomposes under the restriction functor Res_H^K . If the trivial representation of H appears in the decomposition, then this representation is of class one. We have the following classification result:

Theorem A.2. *For each almost effective spherical pair (K, H) , Table 1 gives the classification of all complex irreducible class one representations ρ .*

The multiplicity of the trivial representation of H in $\text{Res}_H^K(\rho)$ is equal to 1 except for the pair $(Sp(n), Sp(n-1))$ where it is $a+1$ if ρ has the highest weight $a\varpi_1 + b\varpi_2$.

In the table the numbers a and b are non-negative integers and k is an integer. In the pair $(Sp(n) \times U(1), Sp(n-1) \times U(1)_m)$, a , b and k satisfy further restriction, denoted by (S): $a+b \geq 1$, $|m|$ divides k , a and $\frac{k}{|m|}$ have the same parity and $\frac{|k|}{|m|} \leq a$. For other pairs, the restrictions are specified in the table.

In Table 1, if $n = 1$ for the $Sp(n)$ factor, then there does not exist $b\varpi_2$ in ρ , i.e., $b = 0$.

The proof of Theorem A.2 is divided into four parts. The results of the first four group pairs in Table 1 are direct consequences of the classical branching rules for those pairs. The

K	H	ρ		n
$SO(n)$	$SO(n-1)$	$a\varpi_1$	$a \geq 1$	$n \geq 3$
$SU(n)$	$SU(n-1)$	$a\varpi_1 + b\varpi_{n-1}$	$a + b \geq 1$	$n \geq 3$
$U(n)$	$U(n-1)_m$	$ae_1 - be_n + m(a-b)(e_1 + \cdots + e_n)$	$a + b \geq 1$	$n \geq 2$
$Sp(n)$	$Sp(n-1)$	$a\varpi_1 + b\varpi_2$	$a + b \geq 1$	$n \geq 1$
$Sp(n) \times Sp(1)$	$Sp(n-1) \times Sp(1)$	$(a\varpi_1 + b\varpi_2) \otimes a\varpi_1$	$a + b \geq 1$	$n \geq 1$
$Sp(n) \times U(1)$	$Sp(n-1) \times U(1)_m$	$(a\varpi_1 + b\varpi_2) \otimes \phi^k$	(S)	$n \geq 1$
G_2	$SU(3)$	$a\varpi_1$	$a \geq 1$	
$Spin(7)$	G_2	$a\varpi_3$	$a \geq 1$	
$Spin(9)$	$Spin(7)$	$a\varpi_1 + b\varpi_4$	$a + b \geq 1$	
$U(1)$	$\{1\}$	ϕ^k	$k \neq 0$	

TABLE 1. Complex class one representations of spherical pairs

second part includes the pairs $(Sp(n) \times Sp(1), Sp(n-1) \times Sp(1))$ and $(Sp(n) \times U(1), Sp(n-1) \times U(1)_m)$. They will be proved in the Section A.2 by using a branching rule of J. Lepowsky. The third part is covered in Section A.3. We will apply Kostant's Branching Theorem to each pair to obtain the class one representations. The result for the last pair $(U(1), \{1\})$ is clear.

In the last section, Section A.4, we will study some properties of these representations, for example, the type, the kernel and the dimension.

A.1. The Class One Representations and Harmonic Polynomials. In this section we prove Theorem A.1 and provide an explicit characterization of class one representations for the pairs $(SU(n), SU(n-1))$ and $(U(n), U(n-1)_m)$ using harmonic polynomials.

Proof of Theorem A.1 : We consider the separable Hilbert space $\mathcal{H}$ of all the square-integrable complex valued functions defined on $\mathbb{S}^{n-1}$ under the following inner-product

$$\langle f_1, f_2 \rangle = \int_{\mathbb{S}^{n-1}} f_1 \bar{f}_2 d\sigma, \quad \forall f_1, f_2 \in \mathcal{H},$$

where $d\sigma$ is an $SO(n)$ bi-invariant measure on $\mathbb{S}^{n-1}$ with total measure = 1.

Using the $SO(n)$ action on $\mathbb{S}^{n-1}$, $SO(n)$ acts on $\mathcal{H}$ by $g \star f(x) = f(g^{-1}x)$ for any $g \in SO(n)$. This representation is called the *left-regular representation* of $SO(n)$ on $\mathcal{H}$ and it is a unitary representation.

Since $SO(n)/SO(n-1) = \mathbb{S}^{n-1}$ is a homogeneous space, $\mathcal{H}$ is also the induced representation $\mathcal{H} = \text{Ind}_{SO(n-1)}^{SO(n)} \text{Id}$, where Id is the trivial representation of $SO(n-1)$ on $\mathbb{C}$, see for

example, Chapter 9.2 in [Kn]. By the Frobenius reciprocity, for any irreducible representation ρ of $SO(n)$, we have the following multiplicities equality

$$[\text{Ind}_{SO(n-1)}^{SO(n)} \text{Id} : \rho] = [\text{Res}_{SO(n-1)}^{SO(n)} \rho : \text{Id}], \quad \text{or} \quad [\mathcal{H} : \rho] = [\text{Res}_{SO(n-1)}^{SO(n)} \rho : \text{Id}],$$

where $[\rho_1 : \rho_2]$ denotes the multiplicity of the irreducible representation ρ_2 in ρ_1 . By the classical branching rule for the pair $(SO(n), SO(n-1))$, $[\text{Res}_{SO(n-1)}^{SO(n)} \rho : \text{Id}] \neq 0$ if and only if ρ has the highest weight ae_1 where a is a nonnegative integer. On the other hand, for any class one representation ρ , $[\text{Res}_{SO(n-1)}^{SO(n)} \rho : \text{Id}] = 1$, or equivalently $[\mathcal{H} : \rho] = 1$.

The representation with highest weight ae_1 can be realized as the representation of the complex valued homogeneous harmonic polynomials with degree a on $\mathbb{S}^{n-1}$ which is denoted by $\mathcal{H}^a$. Furthermore, $\mathcal{H}^a$ is the complexification of the real valued homogeneous harmonic polynomials with degree a . We have the following orthogonal decomposition

$$\mathcal{H} = \bigoplus_{a=0}^{\infty} \mathcal{H}^a.$$

Since $K/H = \mathbb{S}^{n-1}$, K acts on the Hilbert space $\mathcal{H}$ and $\mathcal{H} = \text{Ind}_H^K \text{Id}$, where Id is the trivial representation of H on $\mathbb{C}$. As a subgroup of $SO(n)$, K acts invariantly on $\mathcal{H}^a$. In general $\mathcal{H}^a$ is not an irreducible representation of K and it is decomposed orthogonally into irreducible summands as follows:

$$\text{Res}_K^{SO(n)} \mathcal{H}^a = \bigoplus_b \mathcal{H}^{a,b},$$

and there are only finite many b 's for each value of a . Hence we have the following orthogonal decomposition of $\mathcal{H}$ into irreducible representations of K :

$$\mathcal{H} = \bigoplus_{a,b} \mathcal{H}^{a,b}.$$

Suppose μ is an irreducible representation of K , then μ is of class one for the pair (K, H) if and only if $[\mathcal{H} : \mu] \neq 0$ by the Frobenius reciprocity applied to the pair (K, H) . Then from the above decomposition and the uniqueness of this decomposition, μ is equivalent to $\mathcal{H}^{a,b}$. $\square$

Remark A.3. The existence and uniqueness of the decomposition of $\mathcal{H}$ follow from Theorem 9.4 and Corollary 9.6 in [Kn].

Next using harmonic polynomials, we construct the representation space of the class one representations $\rho = a\varpi_1 + b\varpi_{n-1}$ for the pair $(K, H) = (SU(n), SU(n-1))$ and $\rho = ae_1 - be_n + m(a-b)(e_1 + \cdots + e_n)$ for the pair $(K, H) = (U(n), U(n-1)_m)$. It is already

discussed for the pair $(SU(n), SU(n-1))$ in [Kn]. In the following, we consider the pair $(U(n), U(n-1)_m)$.

Let $\{z_1, \dots, z_n\}$ be the basis of $\mathbb{C}^n$ and $A \in K$ acts on it by the defining representation $\Phi_m(A) : (z_1, \dots, z_n)^\top \mapsto (\det A)^m A \cdot (z_1, \dots, z_n)^\top$. If $A \in SU(n)$, then $\Phi_m(A)$ is just matrix multiplication. Similarly $\Phi_m(A)$ maps $(\bar{z}_1, \dots, \bar{z}_n)^\top$ to $(\det \bar{A})^m \bar{A} \cdot (\bar{z}_1, \dots, \bar{z}_n)^\top$. Let V be the space of homogeneous polynomials in $z_1, \dots, z_n, \bar{z}_1, \dots, \bar{z}_n$ of degree $a+b$. A acts on V by

$$\Phi_m(A) : P\left(\begin{pmatrix} z_1 \\ \vdots \\ z_n \end{pmatrix}, \begin{pmatrix} \bar{z}_1 \\ \vdots \\ \bar{z}_n \end{pmatrix}\right) \mapsto P((\det A)^{-m} A^{-1} \cdot \begin{pmatrix} z_1 \\ \vdots \\ z_n \end{pmatrix}, (\det \bar{A})^{-m} \bar{A}^{-1} \cdot \begin{pmatrix} \bar{z}_1 \\ \vdots \\ \bar{z}_n \end{pmatrix}).$$

Let $V_{a,b}$ be the subspace of polynomials with degree a in $\bar{z}$ and degree b in z . Clearly K leaves $V_{a,b}$ invariant. The Laplacian operator Δ is a multiple of $\sum_{j=1}^n \frac{\partial^2}{\partial z_j \partial \bar{z}_j}$ and it commutes with the K action, so the subspace $H_{a,b}$ of harmonic polynomials in $V_{a,b}$ is an invariant subspace.

If P is a monomial of the form

$$P(z, \bar{z}) = \bar{z}_1^{k_1} \dots \bar{z}_n^{k_n} z_1^{l_1} \dots z_n^{l_n} \quad \text{with} \quad \sum_{j=1}^n k_j = a \quad \text{and} \quad \sum_{j=1}^n l_j = b,$$

then P is a weight vector of weight $\sum_{j=1}^n (k_j - l_j)e_j + m(a-b)(e_1 + \dots + e_n)$. Hence $\rho = ae_1 - be_n + m(a-b)(e_1 + \dots + e_n)$ is the highest weight of the representation $H_{a,b}$ with the weight vector $\bar{z}_1^a z_n^b$. From Exercise 17 of Chapter IV in [Kn], the dimension of $H_{a,b}$ is $\frac{a+b+n-1}{n-1} \binom{a+n-2}{a} \binom{b+n-2}{n-2}$ which equals to the dimension of the irreducible representation of K with the highest weight ρ , see, for example, Proposition A.20, so $H_{a,b}$ is the representation space as desired.

A.2. The Pairs $(Sp(n) \times Sp(1), Sp(n-1) \times Sp(1))$ and $(Sp(n) \times U(1), Sp(n-1) \times U(1)_m)$. For each pair (K, H) , there is an intermediate group L and we will apply the branching rule successively as $\text{Res}_H^K \rho = \text{Res}_H^L (\text{Res}_L^K \rho)$ for any irreducible representation ρ of K .

The branching rule we will use is the one for the pair $(Sp(n), Sp(n-1) \times Sp(1))$ which is established by J. Lepowsky in [Le]. For each irreducible representation $\mu_1 \otimes \mu_2$ of $Sp(n-1) \times Sp(1)$, let $\mu = b_1 e_1 + \dots + b_{n-1} e_{n-1} + b_n e_n$ be the highest weight, then $b_1 \geq \dots \geq b_{n-1} \geq 0$ and $b_n \geq 0$.

Definition A.4. Let $l, m \in \mathbb{Z}$, $m \geq 1$ and let $q_1, \dots, q_m$ be positive integers. Suppose there are l identical balls, m different boxes and the i -th box can contain at most q_i balls. We define $F_m(l; q_1, \dots, q_m)$ to be the number of ways of putting the l balls into the m boxes.

Theorem A.5 (Lepowsky). *Let ρ, μ be irreducible representations of $Sp(n), Sp(n-1) \times Sp(1)$ with highest weights $\rho = a_1 e_1 + \cdots + a_n e_n$, $\mu = b_1 e_1 + \cdots + b_{n-1} e_{n-1} + b_n e_n$ respectively. Let*

$$\begin{aligned} A_1 &= a_1 - \max(a_2, b_1), \\ A_i &= \min(a_i, b_{i-1}) - \max(a_{i+1}, b_i) \quad (2 \leq i \leq n-1), \\ A_n &= \min(a_n, b_{n-1}) \end{aligned}$$

Then the multiplicity $[\text{Res} \rho : \mu] = 0$ unless $\sum_{i=1}^n (a_i + b_i) \in 2\mathbb{Z}$ and $A_1, \dots, A_{n-1} \geq 0$. Under these conditions,

$$\begin{aligned} [\text{Res} \rho : \mu] &= F_{n-1} \left(\frac{1}{2}(b_n - A_1 + \sum_{i=1}^n A_i); A_2, \dots, A_n \right) \\ (A.6) \quad &- F_{n-1} \left(\frac{1}{2}(-b_n - A_1 + \sum_{i=2}^n A_i) - 1; A_2, \dots, A_n \right). \end{aligned}$$

Proof of the 2nd part of Theorem A.2 : Let $K = Sp(n) \times Sp(1)$ and $H = Sp(n-1) \times Sp(1)$, then $L = Sp(n-1) \times Sp(1) \times Sp(1)$ lies between H and K , and the $Sp(1)$ factor in H is diagonally embedded in $Sp(1) \times Sp(1) \subset L$. Suppose $\rho = \rho_1 \otimes \rho_2$ is a class one representation of the pair (K, H) and ρ_1 has the highest weight $a_1 e_1 + \cdots + a_n e_n$ and ρ_2 has the highest weight $b e_1$. Then $\text{Res}_L^K \rho = (\text{Res}_{Sp(n-1) \times Sp(1)}^{Sp(n)} \rho_1) \otimes \rho_2$. Since the trivial representation appears in $\text{Res}_H^K \rho$, it appears in $(\text{Res}_H^L \mu) \otimes \rho_2$ for some irreducible representation $\mu = \mu_1 \otimes \mu_2$ in $\text{Res}_{Sp(n-1) \times Sp(1)}^{Sp(n)} \rho_1$. Under the restriction from $Sp(n-1) \times Sp(1) \times Sp(1)$ to $Sp(n-1) \times \Delta Sp(1)$, $\mu_1 \otimes \mu_2 \otimes \rho_2$ splits as $\mu_1 \otimes \text{Res}_{\Delta Sp(1)}^{Sp(1) \times Sp(1)} (\mu_2 \otimes \rho_2)$. Therefore μ_1 is a trivial representation of $Sp(n-1)$ and μ_2, ρ_2 have the same dimension, i.e., $\text{Res}_{Sp(n-1) \times Sp(1)}^{Sp(n)} \rho_1$ contains the representation $\text{Id} \otimes \rho_2$ which has the highest weight $b e_n$. From Theorem A.5, we have $A_1 = a_1 - a_2 \geq 0$, $A_2 = -a_3 \geq 0$, $\dots$, $A_{n-1} = -a_n \geq 0$, $b + a_1 + a_2$ is an even number and the multiplicity is equal to

$$p = F_{n-1} \left(\frac{1}{2}(b - (a_1 - a_2)); 0, \dots, 0 \right) - F_{n-1} \left(\frac{1}{2}(-b - (a_1 - a_2)) - 1; 0, \dots, 0 \right).$$

Hence $p \neq 0$ if and only if $b = a_1 - a_2$ and in this case we have $p = 1$.

For the second part, we give a proof when $m = 1$ and the general case easily follows from this case. Let $K = Sp(n) \times U(1)$, $H = Sp(n-1) \times U(1)$, $K_1 = Sp(n-1) \times Sp(1) \times U(1)$ and $K_2 = Sp(n-1) \times U(1) \times U(1)$, then we have the embedding $H \subset K_1 \subset K_2 \subset K$ and $U(1)$ in H lies diagonally in $U(1) \times U(1) \subset K_1$. Suppose $\rho = \mu \otimes \phi^k$ is a class one representation of the pair (K, H) and μ has the highest weight $\mu = a_1 e_1 + \cdots + a_n e_n$. From the first part of the proof, we have $a_3 = \cdots = a_n = 0$ and the multiplicity of the trivial representation in

$\text{Res}_H^K \rho$ is equal to the multiplicity of the trivial representation in $(\text{Res}_{U(1)}^{Sp(1)}(a_1 - a_2)e_n) \otimes \phi^k$. Therefore $k, a_1 - a_2$ have the same parity, $|k| \leq a_1 - a_2$ and the multiplicity is 1. $\square$

A.3. The Pairs $(G_2, SU(3))$, $(Spin(7), G_2)$ and $(Spin(9), Spin(7))$. In this section, we will develop the branching rule for each group pair and then prove the corresponding result in Theorem A.2. The main tool is Kostant's Branching Theorem. We quote the statement of the theorem from [Kn].

First we establish the notation we will use. Let K be a connected compact Lie group and let H be a connected closed subgroup. Choose a maximal torus $S \subset H$. The special assumption is that the centralizer of S in K is a maximal torus T of K . Let Δ_K be the set of the roots of $(\mathfrak{k}^\mathbb{C}, \mathfrak{t}^\mathbb{C})$, let Δ_H be the set of roots of $(\mathfrak{h}^\mathbb{C}, \mathfrak{s}^\mathbb{C})$, and let W_K be the Weyl group of Δ_K . Introduce compatible positive systems Δ_K^+ and Δ_H^+ , let bar denote restriction from the dual $(\mathfrak{t}^\mathbb{C})^*$ to the dual $(\mathfrak{s}^\mathbb{C})^*$, and let δ_K be half the sum of the members in Δ_K^+ . The restrictions to $\mathfrak{s}^\mathbb{C}$ of the members of Δ_K^+ , repeated according to their multiplicities, are the nonzero positive weights of $\mathfrak{s}^\mathbb{C}$ in $\mathfrak{k}^\mathbb{C}$. Deleting from this set the members of Δ_H^+ , each with multiplicity 1, we obtain the set Σ of positive weights of $\mathfrak{s}^\mathbb{C}$ in $\mathfrak{k}^\mathbb{C}/\mathfrak{h}^\mathbb{C}$, repeated according to multiplicities. The associated Kostant partition function is defined as follows: $\mathcal{P}(\nu)$ is the number of ways that a member of $(\mathfrak{s}^\mathbb{C})^*$ can be written as a sum of members of Σ , with the multiple versions of a member of Σ being regarded as distinct.

Theorem A.7 (Kostant's Branching Theorem). *Let K be a compact connected Lie group, let H be a closed connected subgroup, suppose that the centralizer in K of a maximal torus S of H is abelian and is therefore a maximal torus T of K , and let other notation be as above. Let ρ be an irreducible representation of K with the highest weight $\rho \in (\mathfrak{t}^\mathbb{C})^*$, and let μ be an irreducible representation of H with the highest weight $\mu \in (\mathfrak{s}^\mathbb{C})^*$. Then the multiplicity of μ in the restriction of ρ to H is given by*

$$[\rho : \mu] = \sum_{w \in W_K} \varepsilon(w) \mathcal{P}(\overline{w(\rho + \delta_K) - \delta_K - \mu}).$$

Remark A.8. To apply this theorem to our special examples, we will use an equivalent assumption on the Cartan subalgebras instead of the maximal tori: the centralizer of $\mathfrak{s}$ in $\mathfrak{h}$ is a Cartan subalgebra $\mathfrak{t}$ of $\mathfrak{k}$. In our three group pairs, this assumption is verified.

To apply Kostant Branching Theorem, we need to work out the explicit embeddings of the Lie algebras. These groups and Lie algebras are well studied, see, for examples, [GIWZ] and [Mu] and references given there.

We start with the pair $(Spin(9), Spin(7))$. Let $\mathbb{O}$ be the set of the Cayley numbers which is isomorphic to $\mathbb{R}^8$ as vector spaces. $SO(8)$ acts on $\mathbb{O}$ by left multiplication and we have the *Principle of Triality*:

Proposition A.9. *For each $\theta_1 \in SO(8)$, there exists $\theta_2, \theta_3 \in SO(8)$ such that*

$$(A.10) \quad \theta_1(x)\theta_2(y) = \theta_3(xy), \quad \text{for any } x, y \in \mathbb{O}$$

Moreover if θ'_2, θ'_3 satisfy the above equation as θ_2, θ_3 , then $(\theta'_2, \theta'_3) = \pm(\theta_2, \theta_3)$.

The infinitesimal version of the above principle is given as follows:

Proposition A.11. *For any $X \in \mathfrak{so}(8)$, there exist $Y, Z \in \mathfrak{so}(8)$ such that*

$$(A.12) \quad (Xx)y + x(Yy) = Z(xy) \quad \text{for any } x, y \in \mathbb{O}.$$

Moreover Y, Z is uniquely determined by X with $Y = \lambda(X), Z = \lambda\kappa(X)$ and λ, κ are two outer automorphisms of $\mathfrak{so}(8)$ with $\lambda^3 = 1, \kappa^2 = 1$ and $\kappa\lambda^2 = \lambda\kappa$.

We identify $\mathbb{O} \oplus \mathbb{O}$ with $\mathbb{R}^{16}$ and then $Spin(9) \subset SO(16)$ acts transitively on $\mathbb{S}^{15} \subset \mathbb{O} \oplus \mathbb{O}$. Let $v_0 \in \mathbb{R} \subset \mathbb{O}$ be the unit length vector, then $Spin(7)$ is the isotropy subgroup at $(v_0, 0)$. Consider the following Hopf fibration:

$$\begin{array}{ccccc} \mathbb{S}^7 & \longrightarrow & \mathbb{S}^{15} & \longrightarrow & \mathbb{S}^8 = \mathbb{O} \cup \{\infty\}, \\ & & (x, y) & \mapsto & y^{-1}\bar{x}. \end{array}$$

The isotropy subgroup of the fiber $\{(x, 0) | x \in \mathbb{O}\}$ is given as follows:

$$(A.13) \quad Spin(8) = \left\{ (\theta_1, \theta_2) \in SO(8) \times SO(8) \mid \exists \theta_3 \in SO(8) \text{ such that } \theta_1(x)\theta_2(y) = \overline{\theta_3(\overline{xy})} \right\},$$

and the embedding of $Spin(7) \subset Spin(8)$ is

$$(A.14) \quad Spin(7) = \left\{ (\theta_1, \theta_2) \in Spin(8) \mid \theta_1 \in SO(7) \subset SO(8) \text{ or } \theta_1(x)\theta_2(y) = \overline{\theta_2(\overline{xy})} \right\}.$$

The automorphism $\theta \mapsto (x \mapsto \overline{\theta(\bar{x})})$ of $SO(8)$ induces the automorphism κ . Let $\theta_1 = \exp(X)$, $\theta_2 = \exp(Y)$ and $\theta_3 = \exp(Z)$. From Proposition A.11, we have $X = \lambda(\kappa Z)$ and $Y = \lambda^2(\kappa Z)$, so we can write the Lie algebra of $Spin(8)$ as

$$\text{Lie}(Spin(8)) = \{(\lambda(X), \lambda^2(X)) \mid X \in \mathfrak{so}(8)\}.$$

From the embedding of $Spin(7) \subset Spin(8)$ in (A.14), since $\theta_1 \in SO(7)$ we have the Lie algebra of $Spin(7)$ as

$$\text{Lie}(Spin(7)) = \{(Y, \lambda(Y)) \mid Y \in \mathfrak{so}(7)\} \subset \text{Lie}(Spin(8)).$$

Therefore if we identify $\text{Lie}(\text{Spin}(8))$ with $\mathfrak{so}(8)$ by its first component, then we have

$$(A.15) \quad \text{Lie}(\text{Spin}(7)) = \{\lambda^2(X) | X \in \mathfrak{so}(7)\}.$$

Let $\{e_i \pm e_j | 1 \leq i < j \leq 4\}$ be the positive root system of $\mathfrak{so}(8)$ with vanishing e_4 on $\mathfrak{so}(7)$, then the automorphism λ^2 induces the following transformation of e_i .

$$(A.16) \quad \lambda^2 \begin{pmatrix} e_1 \\ e_2 \\ e_3 \\ e_4 \end{pmatrix} = \begin{pmatrix} \frac{1}{2} & \frac{1}{2} & \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & \frac{1}{2} & -\frac{1}{2} & -\frac{1}{2} \\ \frac{1}{2} & -\frac{1}{2} & \frac{1}{2} & -\frac{1}{2} \\ -\frac{1}{2} & \frac{1}{2} & \frac{1}{2} & -\frac{1}{2} \end{pmatrix} \begin{pmatrix} e_1 \\ e_2 \\ e_3 \\ e_4 \end{pmatrix}.$$

In (A.14) the θ_2 -component acts transitively on $\mathbb{S}^7 \subset \mathbb{O}$ and the isotropy subgroup at v_0 is G_2 . Since κ is the identity map on $SO(7)$, $\theta_1(x)\theta_2(y) = \theta_2(xy)$ for $\theta_2 \in G_2 \subset SO(7)$ which implies $\theta_1 = \theta_2$. Hence we have

$$G_2 = \{(\theta, \theta) | \theta \in SO(7) \text{ and } \theta(x)\theta(y) = \theta(xy), \text{ for any } x, y \in \mathbb{O}\},$$

and the Lie algebra is

$$\mathfrak{g}_2 = \{(X, X) | X \in \mathfrak{so}(7), X = \lambda(X)\}.$$

If we identify $\text{Lie}(\text{Spin}(7))$ with $\mathfrak{so}(7)$, then we have

$$\mathfrak{g}_2 = \{X | X \in \mathfrak{so}(7), X = \lambda(X)\}.$$

Choose a basis of $\mathbb{O}$ over $\mathbb{R}$, then we can write down the explicit embedding of $\mathfrak{g}_2$, see also in [Mu]. The typical element of $\mathfrak{g}_2$ has the following form:

$$X = \begin{pmatrix} 0 & x_1 - y_1 & x_3 + y_3 & -x_2 + y_2 & -x_4 - y_4 & x_6 + y_6 & x_5 - y_5 \\ -x_1 + y_1 & 0 & a & y_5 & y_6 & y_4 & y_2 \\ -x_3 - y_3 & -a & 0 & x_6 & x_5 & x_2 & x_4 \\ x_2 - y_2 & -y_5 & -x_6 & 0 & b & y_3 & y_1 \\ x_4 + y_4 & -y_6 & -x_5 & -b & 0 & x_1 & x_3 \\ -x_6 - y_6 & -y_4 & -x_2 & -y_3 & -x_1 & 0 & a + b \\ -x_5 + y_5 & -y_2 & -x_4 & -y_1 & -x_3 & -a - b & 0 \end{pmatrix},$$

where $a, b, x_1, \dots, x_6, y_1, \dots, y_6 \in \mathbb{R}$.

The linear functionals $\{e_1, e_2, e_3\}$ of the Cartan subalgebra of $\mathfrak{so}(7)$ satisfy the relation $e_1 = e_2 + e_3$ when restricted to the Cartan subalgebra of $\mathfrak{g}_2$. Suppose $\{\alpha_1, \alpha_2\}$ be the set of the positive simple roots of G_2 where α_1 is the short one, then under the restriction, $\overline{e_3} = \alpha_1$ and $\overline{e_2 - e_3} = \alpha_2$.

We consider the last pair $(G_2, SU(3))$. G_2 acts transitively on the unit sphere $\mathbb{S}^6 = \{x \in \mathbb{O} | \|x\| = 1, \langle x, v_0 \rangle = 0\}$. Let v_1 be a unit element which is orthogonal to v_0 , then the

isotropy group at v_1 is isomorphic to $SU(3) \subset SO(6)$. Since G_2 and $SU(3)$ share the same maximal torus, the restriction of the roots of $\mathfrak{g}_2$ is the identity map.

In the proof we will use the classical branching rules for the special orthogonal groups which are well-known and the proof can be found, for examples, in [Kn].

Theorem A.17 (Branching Rule for $(\mathfrak{so}(2k+1), \mathfrak{so}(2k))$). *The irreducible representation with highest weight $a_1e_1 + a_2e_2 + \cdots + a_ke_k$ of $\mathfrak{so}(2k+1)$ decomposes with multiplicity 1 under $\mathfrak{so}(2k)$, and the representations of $\mathfrak{so}(2k)$ that appear are exactly those with highest weights $\mu = c_1e_1 + c_2e_2 + \cdots + c_ke_k$ such that*

$$a_1 \geq c_1 \geq a_2 \geq c_2 \geq \cdots \geq a_k \geq |c_k|,$$

where $a_i (i = 1, 2, \dots, k)$ are integers or all half integers and $c_j (j = 1, 2, \dots, k)$ are all integers or all half integers.

Theorem A.18 (Branching Rule for $(\mathfrak{so}(2k), \mathfrak{so}(2k-1))$). *The irreducible representation with highest weight $a_1e_1 + a_2e_2 + \cdots + a_ke_k$ of $\mathfrak{so}(2k)$ decomposes with multiplicity 1 under $\mathfrak{so}(2k-1)$, and the representations of $\mathfrak{so}(2k-1)$ that appear are exactly those with highest weights $\mu = c_1e_1 + c_2e_2 + \cdots + c_{k-1}e_{k-1}$ such that*

$$a_1 \geq c_1 \geq a_2 \geq c_2 \geq \cdots \geq c_{k-1} \geq |a_k|,$$

where $a_i (i = 1, 2, \dots, k)$ are integers or all half integers and $c_j (j = 1, 2, \dots, k-1)$ are all integers or all half integers.

Proof of the 3rd Part of Theorem A.2: First let $K = Spin(9)$, $H = Spin(7)$ and $\mathfrak{k} = \text{Lie}(Spin(9))$, $\mathfrak{h} = \text{Lie}(Spin(7))$ be their Lie algebras. $L = Spin(8)$ lies between them and denote its Lie algebra $\text{Lie}(Spin(8))$ by $\mathfrak{l}$. Let

$$\Delta_K^+ = \{e_i \pm e_j | 1 \leq i < j \leq 4\} \cup \{e_i | 1 \leq i \leq 4\},$$

be the positive root system of $\mathfrak{k}$, and then $\mathfrak{l}$ has the positive roots system

$$\Delta_L^+ = \{e_i \pm e_j | 1 \leq i < j \leq 4\}.$$

Since $\mathfrak{k}$ and $\mathfrak{l}$ share the same Cartan subalgebra, $\Sigma = \{e_1, e_2, e_3, e_4\}$. Therefore the branching rule for the pair $(\mathfrak{k}, \mathfrak{l})$ is the same as the classical one for $(\mathfrak{so}(9), \mathfrak{so}(8))$.

Let $f_i = \lambda^2(e_i)$ for $i = 1, 2, 3, 4$. Since λ is an automorphism of $\mathfrak{so}(8)$, we can write

$$\Delta_L^+ = \{f_i \pm f_j | 1 \leq i < j \leq 4\},$$

and $\mathfrak{h}$ has the positive root system

$$\Delta_H^+ = \{f_i \pm f_j | 1 \leq i < j \leq 3\} \cup \{f_i | 1 \leq i \leq 3\}.$$

Therefore $\Sigma = \{f_1, f_2, f_3\}$. The Weyl group W_K also acts on f_i 's and $\delta_K = 3e_1 + 2e_2 + e_3 = 3f_1 + 2f_2 + f_3$. Suppose μ be an irreducible representation of K with the highest weight $\mu = b_1f_1 + b_2f_2 + b_3f_3 + b_4f_4$, then by the classical branching rule for $(\mathfrak{so}(8), \mathfrak{so}(7))$, μ decomposes with multiplicity 1 when restricted to $\mathfrak{h}$, and the representations of $\mathfrak{h}$ that appear are exactly those with highest weights $\tau = c_1f_1 + c_2f_2 + c_3f_3$ such that

$$b_1 \geq c_1 \geq b_2 \geq c_2 \geq b_3 \geq c_3 \geq |b_4|,$$

where $b_i (i = 1, 2, 3, 4)$ and $c_j (j = 1, 2, 3)$ are all integers or half integers. Therefore the class one representations of the pair (K, H) are exactly those with the highest weights bf_1 where $b \in \mathbb{Z}$, or $\frac{1}{2}b(e_1 + \cdots + e_4)$ and the multiplicity of trivial representation is 1.

Now suppose that μ is a class one representation of the pair (K, H) with the highest weight $\rho = a_1e_1 + \cdots + a_4e_4$, then $\text{Res}_L^K \rho$ contains at least one representation with highest weight $\frac{1}{2}b(e_1 + \cdots + e_4)$. From the branching rule of $(\mathfrak{so}(9), \mathfrak{so}(8))$, we have

$$a_1 \geq \frac{1}{2}b \geq a_2 \geq \frac{1}{2}b \geq a_3 \geq \frac{1}{2}b \geq a_4 \geq \frac{1}{2}b.$$

Therefore $a_1 \geq a_2 = a_3 = a_4 \geq 0$ and all are integers or half integers. If a_i 's satisfy the conditions, then ρ contains only one representation with the highest weight μ which implies that $[\text{Res}_H^K \rho : \text{Id}] = 1$.

Next we look at the pair $(K, H) = (\text{Spin}(7), G_2)$. We identify $\mathfrak{k}$ with $\mathfrak{so}(7)$ so that

$$\Delta_K^+ = \{e_i \pm e_j | 1 \leq i < j \leq 3\} \cup \{e_i | 1 \leq i \leq 3\}.$$

By the relation $e_1 = e_2 + e_3$ and the restriction $\overline{e_3} = \alpha_1$, $\overline{e_2 - e_3} = \alpha_2$, we have

$$\overline{\Delta_K^+} = \{\alpha_1, 3\alpha_1 + 2\alpha_2, \alpha_1 + \alpha_2, 3\alpha_1 + \alpha_2, \alpha_2, 2\alpha_1 + \alpha_2, 2\alpha_1 + \alpha_2, \alpha_1 + \alpha_2, \alpha_1\}$$

under the restriction. In terms of α_1, α_2 , $\mathfrak{h}$ has the following positive root system

$$\Delta_H^+ = \{\alpha_1, \alpha_2, \alpha_1 + \alpha_2, 2\alpha_1 + \alpha_2, 3\alpha_1 + \alpha_2, 3\alpha_1 + 2\alpha_2\}.$$

Therefore $\Sigma = \{\alpha_1, \alpha_1 + \alpha_2, 2\alpha_1 + \alpha_2\}$. The Weyl group W_K permutes $\{e_1, e_2, e_3\}$ and changes the sign of each e_i .

Let ρ be a class one representation with the highest weight $\rho = a_1e_1 + a_2e_2 + a_3e_3$ ($a_1 \geq a_2 \geq a_3 \geq 0$). Then from the Kostant's Branching Theorem A.7, by a computation, the multiplicity of the trivial representation Id of G_2 in $\text{Res} \rho$ is

$$p = (a_1 + a_3 + 1) - (a_1 - a_3) - (a_1 + a_3) + \max\{0, a_1 - a_3 - 1\} - (a_1 - a_3 - 1) + \max\{0, a_1 - a_3 - 1\}.$$

Therefore $p \neq 0$ if and only if $a_1 = a_3$, i.e., $a_1 = a_2 = a_3$. Thus in this case we have $p = 1$.

Finally we look at the pair $(K, H) = (G_2, SU(3))$. $\mathfrak{k} = \mathfrak{g}_2$ and $\mathfrak{h} = \mathfrak{su}(3)$ share the same Cartan subalgebra and hence the restriction of the roots is the identity map. $\mathfrak{k}$ has the following positive root system

$$\Delta_K^+ = \{\alpha_1, 3\alpha_1 + \alpha_2, 2\alpha_1 + \alpha_2, 3\alpha_1 + 2\alpha_2, \alpha_1 + \alpha_2, \alpha_2\},$$

where $\alpha_1 = e_3$, $\alpha_2 = e_2 - e_3$ and $e_2 + e_3 - e_1 = 0$. Among the 12 roots of $\mathfrak{k}$, the six long roots consist the root system of $\mathfrak{h}$, i.e.

$$\Delta_H^+ = \{3\alpha_1 + 2\alpha_2, 3\alpha_1 + \alpha_2, \alpha_2\}.$$

Therefore $\Sigma = \{\alpha_1, \alpha_1 + \alpha_2, 2\alpha_1 + \alpha_2\}$. The Weyl group W_K is the symmetry group of the regular hexagon $D_6 = \langle \sigma, \tau \mid \sigma^6 = 1, \tau^2 = 1 \rangle$ with the following operations on α_1, α_2 :

$$\begin{aligned} \sigma &: \alpha_1 \mapsto -\alpha_1 - \alpha_2, & \alpha_2 &\mapsto 3\alpha_1 + 2\alpha_2, \\ \tau &: \alpha_1 \mapsto -\alpha_1 - \alpha_2, & \alpha_2 &\mapsto \alpha_2. \end{aligned}$$

Suppose ρ is a class one representation with the highest weight $\rho = a_1\varpi_1 + a_2\varpi_2 = (2a_1 + 3a_2)\alpha_1 + (a_1 + 2a_2)\alpha_2$, where a_1, a_2 are two nonnegative integers. A computation using Kostant Branching Theorem shows that $a_2 = 0$ and the multiplicity $[\text{Res } \rho : \text{Id}]$ is 1. $\square$

A.4. Properties of Class One Representations. First we consider the types and non-trivial kernels of class one representations. They follow easily from Proposition 23.13 and propositions in §26.3 in [FH].

Proposition A.19. *The type and non-trivial kernel Z of each complex class one representation is listed in Table 2.*

In Table 2, for $K = U(n)(a = b)$ and $K = Sp(n) \times U(1)(k = 0)$, the $\mathbb{Z}_0$ factor in the non-trivial kernel should be interpreted as the group $U(1) = \{z \cdot \text{Id} \mid z \in U(1)\}$. Furthermore, $\mathbb{Z}_2 \subset K$ is generated by $-\text{Id}$ and $\mathbb{Z}_l \subset SU(n)$ (or $U(n)$) is generated by $\varphi_l \cdot \text{Id}$ with $\varphi_l = \exp(\frac{2\pi\sqrt{-1}}{l})$. The notation $\gcd(p, q)$ stands for the greatest common divisor of p, q .

Next we give the dimensions of the class one representations, see [FH].

Proposition A.20. *The dimension of each complex class one representation for the pair (K, H) is listed as follows:*

- $(SO(n), SO(n-1)) : \rho = a\varpi_1,$
 $\dim \rho = \frac{2a+n-2}{a+n-2} \binom{a+n-2}{a}$

K	ρ	Type	Z
$SO(n)$	$a\varpi_1$	real	$\mathbb{Z}_2$: if both n and a are even
$SU(n)$	$a\varpi_1 + b\varpi_{n-1}$	real: if $a = b$ complex: otherwise	$\mathbb{Z}_l$, $l = \gcd(a + b, n)$
$U(n)$	$ae_1 - be_n + m(a - b)(e_1 + \cdots + e_n)$	real: if $a = b$ complex: otherwise	$\mathbb{Z}_{ (a-b)(1+mn) }$: if $a \neq b$
$Sp(n)$	$a\varpi_1 + b\varpi_2$	real: a is even quaternionic: otherwise	$\mathbb{Z}_2$: if a is even
$Sp(n) \times Sp(1)$	$(a\varpi_1 + b\varpi_2) \otimes a\varpi_1$	real	$\mathbb{Z}_2 \times \mathbb{Z}_2$: if $a \neq 0$ is even $\mathbb{Z}_2 \times Sp(1)$: if $a = 0$
$Sp(n) \times U(1)$	$(a\varpi_1 + b\varpi_2) \otimes \phi^k$	complex: $k \neq 0$ real: $k = 0$ and a is even quaternionic: otherwise	$\text{Id} \times \mathbb{Z}_{ k }$: if a is odd $\mathbb{Z}_2 \times \mathbb{Z}_{ k }$: if a is even
G_2	$a\varpi_1$	real	-----
$Spin(7)$	$a\varpi_3$	real	$\mathbb{Z}_2$: if a is even
$Spin(9)$	$a\varpi_1 + b\varpi_4$	real	$\mathbb{Z}_2$: if b is even
$U(1)$	ϕ^k	complex	$\mathbb{Z}_{ k }$: if $ k > 1$

TABLE 2. Type and non-trivial kernel of class one representations

- $(SU(n), SU(n-1)) : \rho = a\varpi_1 + b\varpi_{n-1}$,
 $\dim \rho = \frac{a+b+n-1}{n-1} \binom{a+n-2}{a} \binom{b+n-2}{b}$
- $(U(n), U(n-1)_m) : \rho = ae_1 - be_n + m(a-b)(e_1 + \cdots + e_n)$,
 $\dim \rho = \frac{a+b+n-1}{n-1} \binom{a+n-2}{a} \binom{b+n-2}{b}$
- $(Sp(n), Sp(n-1)) : \rho = a\varpi_1 + b\varpi_2$,
 $\dim \rho = \frac{(a+1)(a+2b+2n-1)}{(a+b+1)(a+b+2n-1)} \binom{a+b+2n-1}{a+b} \binom{b+2n-3}{b}$
- $(Sp(n) \times Sp(1), Sp(n-1) \times Sp(1)) : \rho = (a\varpi_1 + b\varpi_2) \otimes a\varpi_1$,
 $\dim \rho = \frac{(a+1)^2(a+2b+2n-1)}{(a+b+1)(a+b+2n-1)} \binom{a+b+2n-1}{a+b} \binom{b+2n-3}{b}$
- $(Sp(n) \times U(1), Sp(n-1) \times U(1)_m) : \rho = (a\varpi_1 + b\varpi_2) \otimes \phi^k$,
 $\dim \rho = \frac{(a+1)(a+2b+2n-1)}{(a+b+1)(a+b+2n-1)} \binom{a+b+2n-1}{a+b} \binom{b+2n-3}{b}$

- $(G_2, SU(3)) : \rho = a\varpi_1,$
 $\dim \rho = \frac{1}{120}(a+1)(a+2)(a+3)(a+4)(2a+5)$
- $(Spin(7), G_2) : \rho = a\varpi_3,$
 $\dim \rho = \frac{1}{360}(a+1)(a+2)(a+3)^2(a+4)(a+5)$
- $(Spin(9), Spin(7)) : \rho = a\varpi_1 + b\varpi_4,$
 $\dim \rho = \frac{1}{1814400}(a+1)(a+2)(a+3)(b+1)(b+2)(b+3)^2(b+4)(b+5)(a+b+4)(a+b+5)(a+b+6)(2a+b+7)$
- $(U(1), \{1\}) : \rho = \phi^k, \dim \rho = 1.$

Finally for each spherical pair (K, H) , we consider the embedding $\rho : K \longrightarrow SO(l), U(l)$ and $Sp(l)$. We compare the dimension of the sphere K/H and l .

Proposition A.21. *Table 3 is the list of the dimension s of the sphere K/H and the embedding $\rho : K \longrightarrow SO(l)$ with $l \leq k_0(s+1)$ where k_0 is equal to 1, 2 or 4 if ρ is of real, complex or quaternionic type.*

For the complex representations, we have

Proposition A.22. *Table 4 is the list of the dimension s of the sphere K/H and the embedding $\rho : K \longrightarrow U(l)$ with $2l \leq s+1$.*

A similar result for the quaternionic representations is

Proposition A.23. *The quaternionic class one representations $\rho : K \longrightarrow Sp(l)$ with $4l \leq \dim(K/H) + 1$ are exactly the representation $\rho = \varpi_1$ for the pair $(Sp(n), Sp(n-1)) (n \geq 1)$.*

REFERENCES

- [AA] A. V. Alekseevsky and D. V. Alekseevsky, *G-manifolds with one dimensional orbit space*, Ad. in Sov. Math. **8**(1992), 1–31.
- [Be1] A. L. Besse, *Einstein manifolds*, Ergebnisse der Mathematik und ihrer Grenzgebiete (**3**) [Results in Mathematics and Related Areas (**3**)], **10**. Springer-Verlag, Berlin, 1987.
- [Be2] A. L. Besse, *Manifolds all of whose geodesics are closed*, with appendices by D. B. A. Epstein, J. P. Bourguignon, L. Bérard-Bergery, M. Berger and J. L. Kazdan. Ergebnisse der Mathematik und ihrer Grenzgebiete [Results in Mathematics and Related Areas], **93**. Springer-Verlag, Berlin-New York, 1978.
- [Ch] J. Cheeger, *Some examples of manifolds of nonnegative curvature*, J. Diff. Geom. **8**(1973), 623–628.

K	H	ρ	l	$s + 1$	$k_0(s + 1)$
$SO(n)$	$SO(n - 1)$	ϖ_1	n	n	
$SU(n)$	$SU(n - 1)$	$[\varpi_1]_{\mathbb{R}}$	$2n$	$2n$	$4n$
$U(n)$	$U(n - 1)_m$	$[e_1 + m(e_1 + \cdots + e_n)]_{\mathbb{R}}$	$2n$	$2n$	$4n$
$Sp(n)$	$Sp(n - 1)$	$[\varpi_1]_{\mathbb{R}}$	$4n$	$4n$	$16n$
$Sp(n) \times Sp(1)$	$Sp(n - 1) \times Sp(1)$	$\varpi_1 \otimes \varpi_1$	$4n$	$4n$	
$Sp(n) \times U(1)$	$Sp(n - 1) \times U(1)_m$	$[\varpi_1 \otimes \phi^k]_{\mathbb{R}}$	$4n$	$4n$	$8n$ or $16n$
G_2	$SU(3)$	ϖ_1	7	7	
$Spin(7)$	G_2	ϖ_3	8	8	
$Spin(9)$	$Spin(7)$	ϖ_1, ϖ_4	9, 16	16	
$U(2)$	$U(1)_m$	$e_1 - e_2$	3	4	
$Sp(2)$	$Sp(1)$	ϖ_2	5	8	
$Sp(1)$	$\{1\}$	$2\varpi_1$	3	4	
$U(1)$	$\{1\}$	$[\phi^k]_{\mathbb{R}}$	2	2	4
$U(2)$	$U(1)_m$	$[2e_1 + 2m(e_1 + e_2)]_{\mathbb{R}}$ $[3e_1 + 3m(e_1 + e_2)]_{\mathbb{R}}$	6 8	4	8
$U(3)$	$U(2)_m$	$[2e_1 + 2m(e_1 + e_2 + e_3)]_{\mathbb{R}}$	12	6	12
$SU(3)$	$SU(2)$	$[2\varpi_1]_{\mathbb{R}}$	12	6	12
$Sp(2)$	$Sp(1)$	$[\varpi_1 + \varpi_2]_{\mathbb{R}}$	32	8	32
$Sp(1)$	$\{1\}$	$[3\varpi_1]_{\mathbb{R}}, [5\varpi_1]_{\mathbb{R}}, [7\varpi_1]_{\mathbb{R}}$	8, 12, 16	4	16

TABLE 3. Orthogonal Class One Representation with Small Dimension

K	H	ρ	l	$s + 1$
$SU(n)$	$SU(n - 1)$	ϖ_1, ϖ_{n-1}	n, n	$2n$
$U(n)$	$U(n - 1)$	$e_1, -e_n$	n, n	$2n$
$Sp(n)$	$Sp(n - 1)$	ϖ_1	$2n$	$4n$
$Sp(n) \times U(1)$	$Sp(n - 1) \times U(1)_m$	$\varpi_1 \otimes \phi^k$	$2n$	$4n$

TABLE 4. Complex Class One Representation with Small Dimension

[FH] W. Fulton and J. Harris, *Representation theory. A first course*, Graduate Texts in Mathematics, **129**, Springer-Verlag, New York, 1991.

[GIWZ] H. Gluck, F. Warner and W. Ziller, *The geometry of the Hopf fibrations*, Enseign. Math.(2) **32**(1986), 173–198.

- [GWZ] K. Grove, B. Wilking and W. Ziller, *Positively curved cohomogeneity one manifolds and 3-Sasakian geometry*, J. Diff. Geom. **78**(2008), 33–111.
- [GVWZ] K. Grove, L. Verdiani, B. Wilking and W. Ziller, *Non-negative curvature obstructions in cohomogeneity one and the Kervaire spheres*, Ann. del. Scuola Norm. Sup. **5**(2006), 159–170.
- [GVZ] K. Grove, L. Verdiani and W. Ziller, *A new type of a positively curved manifold*, Preprint(2008), arXiv math:08092304.
- [GZ1] K. Grove and W. Ziller, *Curvature and symmetry of Milnor spheres*, Ann. of Math. **152**(2000), 331–367.
- [GZ2] K. Grove and W. Ziller, *Cohomogeneity one manifolds with positive Ricci curvature*, Invent. Math. **149**(2002), 619–646.
- [Ho] C. Hoelscher, *Classification of cohomogeneity one manifolds in low dimensions*, Preprint(2007), arXiv math:07121327.
- [Kn] A. Knapp, *Lie groups beyond an introduction. Second edition*, Progress in Mathematics, **140**, Birkhäuser Boston, Inc., Boston, MA, 2002.
- [Le] J. Lepowsky, *Multiplicity formulas for certain semisimple Lie groups*, Bull. Amer. Math. Soc. **77**(1971), 601–605.
- [Mu] S. Murakami, *Exceptional simple Lie groups and related topics in recent differential geometry*, Differential geometry and topology(Tianjin, 1986–87), 183–221, Lecture Notes in Math., **1369**, Springer, Berlin, 1989.
- [St] E. Straume, *Compact Connected Lie Transformation Groups on Spheres with Low Cohomogeneity, I*, Mem. Amer. Math. Soc. **119**(1996), no. 569.
- [Ve1] L. Verdiani, *Invariant metrics on cohomogeneity one manifolds*, Geom. Dedicata **77**(1999), 77–111.
- [Ve2] L. Verdiani, *Cohomogeneity one Riemannian manifolds of even dimension with strictly positive curvature. I*, Math. Z. **241**(2002), 329–339.
- [Ve3] L. Verdiani, *Cohomogeneity one manifolds of even dimension with strictly positive sectional curvature*, J. Diff. Geom. **68**(2004), 31–72.
- [VK] N. Vilenkin and A. Klimyk, *Representation of Lie groups and special functions*, Vol. I and Vol. II, Mathematics and its Applications(Soviet Series), **72** and **74**, Kluwer Academic Publishers Group, Dordrecht, 1991 and 1993.
- [Wa] N. Wallach, *Minimal immersions of symmetric spaces into spheres*, pp. 1–40. Pure and Appl. Math., Vol. **8**, Dekker, New York, 1972.
- [Wil] B. Wilking, *A duality theorem for Riemannian foliations in nonnegative sectional curvature*, Geom. Funct. Anal. **17**(2007), 1297–1320.
- [Wi2] B. Wilking, *Nonnegatively and positively curved manifolds*, Surv. Differ. Geom. **11**, 25–62, ed. K. Grove and J. Cheeger, Int. Press, Somerville, MA, 2007.
- [Zi] W. Ziller, *Examples of Riemannian manifolds with non-negative sectional curvature*, Surv. Differ. Geom. **11**, 63–102, ed. K. Grove and J. Cheeger, Int. Press, Somerville, MA, 2007.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF PENNSYLVANIA

E-mail address: hech@math.upenn.edu